

FRONT



CONSTRUCTION

BOUND BOOKLET

.375"

4.75"

.375"



**TOUGH ON
APHIDS AND
BEETLES,
GENTLE ON
GARDENS***

GARDEN^{TECH}
Sevin
**INSECT KILLER
CONCENTRATE**
TRUSTED BRAND FOR OVER 50 YEARS

QUICK FACTS

PRODUCT FACTS  Kills by contact Visible results in minutes Allow product to dry completely before people and pets return to the lawn	FRUITS & VEGETABLES  Protects over 250 listed fruits and vegetables Tomato, Pepper, Squash, Cucumber, Broccoli, Kale, Lettuce, Blackberry, Carrot
WHERE TO USE  Fruit and Vegetable Gardens, Ornamental Plants and Shrubs, Flowers and Ground Covers, Around Structures, Lawns	KILLS  Imported Cabbageworm, Cutworms, Armyworms, Flea Beetles, Stink Bugs, Whiteflies, Aphids, Japanese Beetles and other listed lawn & garden insects
HOW MUCH TO USE  4 fl. oz. per gallon of water Treats 8,000 sq. ft. of lawn	

Distributed by:
TechPac, L.L.C.
1000 Parkwood Circle, Suite 700
Atlanta, GA, 30339

EPA Reg. No. 279-3347-71004
EPA Est. 67572-GA-1^a, 67572-GA-2^b
Subscript used is first letter of run code on container.

If you have a medical emergency, please call 1-(800) 420-9347

*When used as directed

Open for Precautionary Statements,
First Aid and Complete Directions for Use



6 13499 01016 2
AAA08026RT


6.0"

5.5"

FOR VISUAL REFERENCE ONLY

BASE LABEL

BASE TEXT (Underneath Book)



FRONT COVER
OF BOOKLET
FRONT COVER
(Page 1)

4.5"

.125"

.125"

5.75"

6.0"

4.75"

.125"

TOUGH ON APHIDS AND BEETLES, GENTLE ON GARDENS*

GARDEN TECH

Sevin
INSECT KILLER
CONCENTRATE

TRUSTED BRAND FOR OVER 50 YEARS

QUICK FACTS

PRODUCT FACTS 	Kills by contact Visible results in minutes Allow product to dry completely before people and pets return to the lawn	FRUITS & VEGETABLES 	Protects over 250 listed fruits and vegetables Tomato, Pepper, Squash, Cucumber, Broccoli, Kale, Lettuce, Blackberry, Carrot
WHERE TO USE 	Fruit and Vegetable Gardens, Ornamental Plants and Shrubs, Flowers and Ground Covers, Around Structures, Lawns	KILLS 	Imported Cabbageworm, Cutworms, Armyworms, Flea Beetles, Stink Bugs, Whiteflies, Aphids, Japanese Beetles and other listed lawn & garden insects
HOW MUCH TO USE 	4 fl. oz. per gallon of water Treats 8,000 sq. ft. of lawn		

Distributed by:
TechPac, L.L.C.
 1000 Parkwood Circle, Suite 700
 Atlanta, GA, 30339

EPA Reg. No. 279-3347-71004
 EPA Est. 67572-GA-1, 67572-GA-2
 Subscript used is first letter of run code on container.

If you have a medical emergency, please call 1-(800) 420-9347

*When used as directed

Open for Precautionary Statements, First Aid and Complete Directions for Use

6 13499 01016 2

AAA080216RT

**NOTE: The total number of pages must be divisible by four (4).
The maximum number of pages for this book is forty (40).**

BOOKLET PAGES

BOOKLET TEXT (Pages 2-40)

.25" ← 4.25" → .25"																																						
.25"	<table border="1"><tr><td>FIRST AID</td><td>FOR OUTDOOR USES - continued</td></tr><tr><td>If swallowed • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person. </td><td>Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications, only.</td></tr><tr><td>If in eyes • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. </td><td>Do not water the treated area to the point of run-off. Do not make applications during rain. Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow this product to enter any drain during or after application.</td></tr><tr><td>HOTLINE NUMBER Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-420-9347 for Emergency Assistance. </td><td>If you have a medical emergency, need consultation on diagnosis, therapy and analysis please call 1-(800) 420-9347. If you have questions relating to product use or application, please call 1-(866) 945-5033. If you have a product spill or transportation emergency call 1-(800) 424-9300. For Consumer Refund: TechPac, LLC PO Box 95437 Palatine, IL 60095-0437</td></tr><tr><td>NOTE TO PHYSICIAN Skin contact (vapor or powder) may be followed by transient tingling or numbness, usually of the face, but this subsides without treatment. This product is a pyrethroid. If large amounts have been ingested, the stomach and intestines should be evacuated. Treatment is symptomatic and supportive. </td><td>IMPORTANT – Directions for Storage and Disposal</td></tr><tr><td>For Information Regarding the Use of this Product Call 1-866-945-5033.</td><td>Storage Keep from freezing. Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place, preferably in a locked storage area and avoid excess heat. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into drink containers.</td></tr><tr><td>PRECAUTIONARY STATEMENTS ENVIRONMENTAL HAZARDS This pesticide is extremely toxic to fish, aquatic invertebrates, oysters and shrimp. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. Care should be used when spraying to avoid fish and reptile pets in/around ornamental ponds. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are foraging the treatment area. </td><td>Pesticide Disposal and Container Handling Nonrefillable container. Do not reuse or refill this container. If empty, place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.</td></tr><tr><td>DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling. SHAKE WELL BEFORE USING</td><td>Dealers Must Sell in Original Packages Only The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. To the extent consistent with applicable law, User agrees to hold TechPac LLC and Seller harmless for any claims relating to such factors. </td></tr><tr><td>USE RESTRICTIONS Do not treat pets with this product. Do not apply this product in a way that will contact adults, children, or pets, either directly or through drift. Do not allow adults, children, or pets to enter the treated area until sprays have dried. Remove pets, birds, and cover fish aquariums before spraying. Avoid contamination of food and feedstuffs. Remove or cover exposed food and drinking water before application. Do not use in food handling establishments in which food is held, processed, prepared, and/or served. When used in the home, remove or cover dishes, utensils, food processing equipment, and food preparation surfaces or wash them before use. Do not apply directly to animals. Do not use in aircraft. </td><td>SPECIFIC USE DIRECTIONS Sevin® Insect Killer Concentrate applied at 4 fl. oz. per gallon of water delivers a spray solution containing 0.011% zeta-cypermethrin. For all applications, this product provides control of each pest as listed in the Pest Table (see page 4). </td></tr><tr><td>FOR OUTDOOR USES All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses: (1) Treatment to soil or vegetation around structures; (2) Applications to lawns, turf, and other vegetation; (3) Applications to building foundations, up to a maximum height of 3 feet. </td><td><table border="1"><tr><td rowspan="6">MEASURING INSTRUCTIONS</td><td>Fl. Oz. of Sevin® Insect Killer Concentrate</td><td>Amount of Water</td></tr><tr><td>1</td><td>1 Quart (32 oz)</td></tr><tr><td>2</td><td>½ Gallon (64 oz)</td></tr><tr><td>4</td><td>1 Gallon (128 oz)</td></tr><tr><td>6</td><td>1½ Gallon (192 oz)</td></tr><tr><td>8</td><td>2 Gallon (256 oz)</td></tr></table> PUMP-STYLE TANK SPRAYERS • Close sprayer tightly and pressurize with hand pump. For spot applications, use a coarse spray nozzle. For larger area applications, use a fan spray nozzle. • Hold tip of wand 12 inches from target area while spraying. DIAL-STYLE HOSE-END SPRAYERS • Fill sprayer jar with enough product for the area to be treated. DO NOT ADD WATER. • Close sprayer, set dial, hook to garden hose and apply. • Pour any unused product back into the product's original container. </td></tr><tr><td> .25" </td><td> 6.0" </td></tr><tr><td colspan="2"> ← 4.75" → </td></tr></table>	FIRST AID	FOR OUTDOOR USES - continued	If swallowed • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.	Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications, only.	If in eyes • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.	Do not water the treated area to the point of run-off. Do not make applications during rain. Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow this product to enter any drain during or after application.	HOTLINE NUMBER Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-420-9347 for Emergency Assistance.	If you have a medical emergency, need consultation on diagnosis, therapy and analysis please call 1-(800) 420-9347. If you have questions relating to product use or application, please call 1-(866) 945-5033. If you have a product spill or transportation emergency call 1-(800) 424-9300. For Consumer Refund: TechPac, LLC PO Box 95437 Palatine, IL 60095-0437	NOTE TO PHYSICIAN Skin contact (vapor or powder) may be followed by transient tingling or numbness, usually of the face, but this subsides without treatment. This product is a pyrethroid. If large amounts have been ingested, the stomach and intestines should be evacuated. Treatment is symptomatic and supportive.	IMPORTANT – Directions for Storage and Disposal	For Information Regarding the Use of this Product Call 1-866-945-5033.	Storage Keep from freezing. Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place, preferably in a locked storage area and avoid excess heat. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into drink containers.	PRECAUTIONARY STATEMENTS ENVIRONMENTAL HAZARDS This pesticide is extremely toxic to fish, aquatic invertebrates, oysters and shrimp. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. Care should be used when spraying to avoid fish and reptile pets in/around ornamental ponds. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are foraging the treatment area.	Pesticide Disposal and Container Handling Nonrefillable container. Do not reuse or refill this container. If empty, place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.	DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling. SHAKE WELL BEFORE USING	Dealers Must Sell in Original Packages Only The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. To the extent consistent with applicable law, User agrees to hold TechPac LLC and Seller harmless for any claims relating to such factors.	USE RESTRICTIONS Do not treat pets with this product. Do not apply this product in a way that will contact adults, children, or pets, either directly or through drift. Do not allow adults, children, or pets to enter the treated area until sprays have dried. Remove pets, birds, and cover fish aquariums before spraying. Avoid contamination of food and feedstuffs. Remove or cover exposed food and drinking water before application. Do not use in food handling establishments in which food is held, processed, prepared, and/or served. When used in the home, remove or cover dishes, utensils, food processing equipment, and food preparation surfaces or wash them before use. Do not apply directly to animals. Do not use in aircraft.	SPECIFIC USE DIRECTIONS Sevin® Insect Killer Concentrate applied at 4 fl. oz. per gallon of water delivers a spray solution containing 0.011% zeta-cypermethrin. For all applications, this product provides control of each pest as listed in the Pest Table (see page 4).	FOR OUTDOOR USES All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses: (1) Treatment to soil or vegetation around structures; (2) Applications to lawns, turf, and other vegetation; (3) Applications to building foundations, up to a maximum height of 3 feet.	<table border="1"><tr><td rowspan="6">MEASURING INSTRUCTIONS</td><td>Fl. Oz. of Sevin® Insect Killer Concentrate</td><td>Amount of Water</td></tr><tr><td>1</td><td>1 Quart (32 oz)</td></tr><tr><td>2</td><td>½ Gallon (64 oz)</td></tr><tr><td>4</td><td>1 Gallon (128 oz)</td></tr><tr><td>6</td><td>1½ Gallon (192 oz)</td></tr><tr><td>8</td><td>2 Gallon (256 oz)</td></tr></table> PUMP-STYLE TANK SPRAYERS • Close sprayer tightly and pressurize with hand pump. For spot applications, use a coarse spray nozzle. For larger area applications, use a fan spray nozzle. • Hold tip of wand 12 inches from target area while spraying. DIAL-STYLE HOSE-END SPRAYERS • Fill sprayer jar with enough product for the area to be treated. DO NOT ADD WATER. • Close sprayer, set dial, hook to garden hose and apply. • Pour any unused product back into the product's original container.	MEASURING INSTRUCTIONS	Fl. Oz. of Sevin® Insect Killer Concentrate	Amount of Water	1	1 Quart (32 oz)	2	½ Gallon (64 oz)	4	1 Gallon (128 oz)	6	1½ Gallon (192 oz)	8	2 Gallon (256 oz)	.25"	6.0"	← 4.75" →	
FIRST AID	FOR OUTDOOR USES - continued																																					
If swallowed • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.	Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications, only.																																					
If in eyes • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.	Do not water the treated area to the point of run-off. Do not make applications during rain. Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow this product to enter any drain during or after application.																																					
HOTLINE NUMBER Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-420-9347 for Emergency Assistance.	If you have a medical emergency, need consultation on diagnosis, therapy and analysis please call 1-(800) 420-9347. If you have questions relating to product use or application, please call 1-(866) 945-5033. If you have a product spill or transportation emergency call 1-(800) 424-9300. For Consumer Refund: TechPac, LLC PO Box 95437 Palatine, IL 60095-0437																																					
NOTE TO PHYSICIAN Skin contact (vapor or powder) may be followed by transient tingling or numbness, usually of the face, but this subsides without treatment. This product is a pyrethroid. If large amounts have been ingested, the stomach and intestines should be evacuated. Treatment is symptomatic and supportive.	IMPORTANT – Directions for Storage and Disposal																																					
For Information Regarding the Use of this Product Call 1-866-945-5033.	Storage Keep from freezing. Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place, preferably in a locked storage area and avoid excess heat. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into drink containers.																																					
PRECAUTIONARY STATEMENTS ENVIRONMENTAL HAZARDS This pesticide is extremely toxic to fish, aquatic invertebrates, oysters and shrimp. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. Care should be used when spraying to avoid fish and reptile pets in/around ornamental ponds. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are foraging the treatment area.	Pesticide Disposal and Container Handling Nonrefillable container. Do not reuse or refill this container. If empty, place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.																																					
DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling. SHAKE WELL BEFORE USING	Dealers Must Sell in Original Packages Only The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. To the extent consistent with applicable law, User agrees to hold TechPac LLC and Seller harmless for any claims relating to such factors.																																					
USE RESTRICTIONS Do not treat pets with this product. Do not apply this product in a way that will contact adults, children, or pets, either directly or through drift. Do not allow adults, children, or pets to enter the treated area until sprays have dried. Remove pets, birds, and cover fish aquariums before spraying. Avoid contamination of food and feedstuffs. Remove or cover exposed food and drinking water before application. Do not use in food handling establishments in which food is held, processed, prepared, and/or served. When used in the home, remove or cover dishes, utensils, food processing equipment, and food preparation surfaces or wash them before use. Do not apply directly to animals. Do not use in aircraft.	SPECIFIC USE DIRECTIONS Sevin® Insect Killer Concentrate applied at 4 fl. oz. per gallon of water delivers a spray solution containing 0.011% zeta-cypermethrin. For all applications, this product provides control of each pest as listed in the Pest Table (see page 4).																																					
FOR OUTDOOR USES All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses: (1) Treatment to soil or vegetation around structures; (2) Applications to lawns, turf, and other vegetation; (3) Applications to building foundations, up to a maximum height of 3 feet.	<table border="1"><tr><td rowspan="6">MEASURING INSTRUCTIONS</td><td>Fl. Oz. of Sevin® Insect Killer Concentrate</td><td>Amount of Water</td></tr><tr><td>1</td><td>1 Quart (32 oz)</td></tr><tr><td>2</td><td>½ Gallon (64 oz)</td></tr><tr><td>4</td><td>1 Gallon (128 oz)</td></tr><tr><td>6</td><td>1½ Gallon (192 oz)</td></tr><tr><td>8</td><td>2 Gallon (256 oz)</td></tr></table> PUMP-STYLE TANK SPRAYERS • Close sprayer tightly and pressurize with hand pump. For spot applications, use a coarse spray nozzle. For larger area applications, use a fan spray nozzle. • Hold tip of wand 12 inches from target area while spraying. DIAL-STYLE HOSE-END SPRAYERS • Fill sprayer jar with enough product for the area to be treated. DO NOT ADD WATER. • Close sprayer, set dial, hook to garden hose and apply. • Pour any unused product back into the product's original container.	MEASURING INSTRUCTIONS	Fl. Oz. of Sevin® Insect Killer Concentrate	Amount of Water	1	1 Quart (32 oz)	2	½ Gallon (64 oz)	4	1 Gallon (128 oz)	6	1½ Gallon (192 oz)	8	2 Gallon (256 oz)																								
MEASURING INSTRUCTIONS	Fl. Oz. of Sevin® Insect Killer Concentrate		Amount of Water																																			
	1		1 Quart (32 oz)																																			
	2		½ Gallon (64 oz)																																			
	4		1 Gallon (128 oz)																																			
	6		1½ Gallon (192 oz)																																			
	8	2 Gallon (256 oz)																																				
.25"	6.0"																																					
← 4.75" →																																						

NOTE: The total number of pages must be divisible by four (4).

The maximum number of pages for this book is forty (40).

BOOKLET PAGES

BOOKLET TEXT (Pages 2-40)

.25"		4.25"		.25"	
.25"					
APPLICATION INSTRUCTIONS		• Apply where insects are expected to appear or when insects or damage are seen. Repeat as necessary according to site-specific directions listed below. • Apply in calm weather. • Begin spraying at the farthest point of the treatment area and work backward to avoid contact with wet treated surfaces. • Spray thoroughly to wet plant surfaces up to the point of runoff. • Sprays must be dry in treated areas before people or pets re-enter. • Do not apply to trees taller than 10 feet. Consider hiring a licensed professional.		VEGETABLES	
		Cress (Garden, Upland (yellow rocket, winter cress)), Dandelion, Dock (sorrel), Endive (escarole), Fennel/Florence (finocchio), Garden Cress, Lettuce (head and leaf), Orach, Parsley, Purslane (garden), Purslane (winter), Radicchio (red chicory), Rhubarb, Spinach (including New Zealand and vine, Malabar spinach, Indian spinach), Swiss Chard		1 Day	
		Peanuts		7 Days	
		Root and Tuber Vegetables (except sugar beets): Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Garden Beet, Edible Burdock, Edible Canna, Carrot, Cassava (Bitter and Sweet), Celeriac (Celery Root), Chayote (Root), Turnip-Rooted Chervil, Chicory, Chufa, Dasheen (Taro), Ginger, Ginseng, Horseradish, Leren, Turnip-Rooted Parsley, Parsnip, Potato, Oriental Radish (Daikon), Radish, Rutabaga, Salsify (Oyster Plant), Black Salsify, Spanish Salsify, Skirret, Sweet Potato, Tanier (Cocoyam), Turmeric, Turnip, Yam Bean, Yam (True)		1 Day	
		Sunflowers: Castor Oil Plant, Chinese Tallowtree, Euphorbia, Evening Primrose, Jojoba, Niger Seed, Rose Hip, Stokes Aster, Tallowwood, Tea Oil Plant, and Vernonia		30 Days	
		BERRIES, FRUITS, AND NUTS			
		Use Sevin® Insect Killer Concentrate as a spot treatment for pest infestations on the berries, fruits and nuts listed below. Applications to trees over 10 feet or trees with areas inaccessible to sprays are not recommended. Apply spray mixture to upper and lower leaves, around fruit clusters, and on branches and trunks up to the point of runoff. Reapply if insects return, but not more than once every seven days and according to pre-harvest interval.			
		BERRIES, FRUITS, AND NUTS			
		Berries Crop Group including: Blackberry, Loganberry, Red and Black Raspberry, Blueberry, Highbush and Lowbush, Currant, Elderberry, Gooseberry, Huckleberry, cultivars and/or hybrids of these			
		1 Day			
		Grapes			
		1 Day			
		Pome Fruit Group including: Apple, Crabapple, Loquat, Mayhaw, Pear, Oriental Pear, and Quince			
		14 Days			
		Stone Fruit Group including: Apricot, Cherry (Sweet and Tart), Nectarine, Peach, Plum (including Chickasaw Plum, Damson Plum, and Japanese Plum), Plumcot, Prune (fresh)			
		14 Days			
		Tree Nuts Group including: Almond, Beech nut, Brazil nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory nut, Macadamia nut, Pecan, Pistachio, Walnut (Black and English)			
		7 Days			
		Tropical Fruits including: Avocado, Black Sapote, Canistel, Maney Sapote, Mango, Papaya, Sapodilla, Star Apple			
		1 Day			
		ORNAMENTAL TREES, SHRUBS, FLOWERS, AND GROUND COVERS			
		Use Sevin® Insect Killer Concentrate as a spot treatment for pest infestations on ornamental trees, shrubs, flowers and ground covers. Spray up to the point of run-off making sure to cover all sides of leaves and trunk. For ground covers, bedding plants, foliage, annuals and perennials, apply as a uniform spray. Reapply if pests return but repeat applications should be limited to no more than once per seven days. Certain plants such as annuals may be sensitive to the final spray solution. A limited area should be treated and observed for one week prior to application to the entire planting.			
		OUTDOOR PERIMETER TREATMENTS			
		Apply to the point of run-off as a residual spray to vegetation such as bushes and trees around buildings and other structures. For ants, apply to any trails, around doors and windows and other places where ants may be found. To prevent pests from entering buildings, apply to a band			
		3			
.25"		4.75"			

**NOTE: The total number of pages must be divisible by four (4).
The maximum number of pages for this book is forty (40).**

BOOKLET PAGES

BOOKLET TEXT (Pages 2-40)

.25"	4.25"	.25"
.25"		
5.5"		6.0"
.25"		
OUTDOOR PERIMETER TREATMENTS - continued of soil and vegetation 5 to 10 feet wide around and adjacent to the building. Also treat the building foundation to a height of 2 to 3 feet where pests are active and may find entrance. Apply as a coarse spray. The surface to be treated should be dry at the time of application. LAWNS Uniformly apply 4 fl. oz. Sevin® Insect Killer Concentrate per gallon of water per 250 sq ft area. For best results, apply after mowing (lawn should not be longer than 3 inches at the time of application). Repeat applications are necessary only if there are signs of renewed insect activity. PESTS CONTROLLED Caterpillars, Butterflies and Moths Tent forming caterpillars, cocoon forming caterpillars, chrysalis forming caterpillars. Leaf feeding caterpillars (including leaves of flowering plants, bushes and other ornamentals, vegetables and trees), azalea leafminers, green cloverworm, fruit eating caterpillars, legume feeding caterpillars, vegetable feeding caterpillars, stem and trunk boring caterpillars, and leafrolling caterpillars. Caterpillars of the Yucca Moth family such as Fairy moths and Yucca moths; Clothes moth family; bagworm moth family, Gelechiid moth family such as Potato tuber moths; Ermine Moth family such as Ailanthus Webworm moth; Clear Winged moth family such as Manroot borer, Squash Vine borer, Doll's Clearwing moth, California Sycamore borer; Tortricid Moth family (millers) such as Fruit Tree Leafroller moth, Orange Tortrix, and Codling moth; Skippers such as Least Skipper, Silver-spotted Skipper, and Long-tail Skipper; Swallowtails such as Green swallowtail, Zebra Swallowtail, Giant Swallowtail (Orange Tail, Orange Puppy), Pale Swallowtail, Eastern Tiger Swallowtail, Eastern Black Swallowtail, Spicebush Swallowtail, and Phoebus; Whites, Sulphurs and Orange Tips family such as Sara Orange Tip, Great Southern White, and European Cabbage Butterfly; Cossamer-Winged Butterflies family such as Great Purple Hairstreak, Spring Azure, Early Hairstreak, Eastern Tailed blue, Harvester, Silvery Blue, Brown Elfin, Marine Blue, American Copper, and Gray Hairstreak; Metalmark family such as Northern Metalmark; Snout Butterflies family such as Common Snout Butterfly; Brush-Footed Butterfly family such as Gulf Fritillary, White Peacock, Harris' Checkerspot, Painted Lady (Thistle butterfly), American Painted Lady (Painted Beauty), Baltimore, Zebra (Yellow-barred Heliconian), Viceroys, White Admiral (Banded Purple), Mourning Cloak, Milbert's Tortoise Shell, Pearl Crescent, Question Mark, Buckeye, Great Spangled Fritillary, and Red Admiral; Hackberry and Goatweed Butterfly family such as Hackberry butterfly; Satyrs, Nymphs and Arctics family such as Wood Nymph (American Grayling), White Wood Satyr, Eyed Brown (Grass Nymph), Pearly Eye, and Chryxus Arctic; Milkweed Butterfly family such as Queen and Monarch; Smoky Moth family such as Grape Leaf Skeletonizer; Slug Caterpillar Moth family such as Saddleback Caterpillar Moth; Pyralid Moths family such as Mediterranean Flour Moth Sod Webworm Moth, Indian Meal Moth, Meal Moth; Plume Moth family such as Artichoke Plume Moth; Measuringworm Moth family such as Tulip Tree Beauty, California Cankerworm Moth, and Large California Spanworm Moth; Tent Caterpillar Moth family such as Western Tent Caterpillar Moth; Giant Silkworm Moth family such as Luna Moth, Polyphemus moth, Io Moth, Prometheus Moth (Spicebush Silkworm), Regal Moth (Royal Walnut Moth), Rosy Maple Moth, Imperial Moth, Sheep Moth, Cecropia Moth, and Cynthia Moth; Sphinx moth family such as Virginia-creeper Sphinx, Pandora Sphinx, Hummingbird Moth (Common Clearwing), White-lined Sphinx, Tomato Hornworm Moth, Tobacco Hornworm Moth, Big Poplar Sphinx, Cerisy's Sphinx, and Wild-cherry Sphinx; Prominent's family such as Tentacled Prominent's (Puss Moths), and Red-humped Appleworm Moth; Tiger Moth family such as Ornate Tiger Moth, Yellow Woolly Bear Moth, Acraea Moth, Milkweed Tiger Moth, Spotted Tiger Moth, Colona, Woolly Bear Caterpillar Moth (Banded Woolly Bear), Lichen Moth, and Rattlebox Moth; Ctenuchid Moth family such as Virginia Ctenuchid Moth; Tussock moth family such as Gypsy Moth, White-marked Tussock Moth, Owllet Moth family such as Eight-spotted Forester, Black Witch, Alfalfa Looper, Sweetheart Underwing, Locust Underwing, and Hebrew PESTS CONTROLLED - continued Common Bristletails Silverfish Family such as Silverfish, and Firebrat Springtails Podurid Springtail family such as Snow Pea, Seashore Springtail; Entomobryid Springtail family such as Ainsley's Springtail; Globular Springtail family Cockroaches Blattid Cockroach Family such as Oriental Cockroach (Asiatic Cockroach, Shad Cockroach), and American Cockroach (Waterbug); Blattellid Cockroaches family such as German Cockroach (Croton Bug) Termites Rotting-wood termite family such as Pacific Coast Termite; Subterranean termite family such as subterranean termites. Not intended to replace structural treatments. For termites – not recommended as sole protection against termites. Such applications are not a substitute for mechanical alterations, soil or foundation treatment but merely as a supplement. For active infestations, get a professional inspection. Earwigs Long-Horned Earwig family such as Ring-legged Earwig, and Riparian Earwig; Common Earwig family such as European Earwig Grasshoppers & Crickets Pygmy Grasshopper family such as Aztec Pygmy Grasshopper; Short-Horned Grasshopper family such as Creosote Bush Grasshopper, Lubber Grasshopper, Painted Grasshopper, Carolina Locust, Dragon Lubber Grasshopper (Gray Dragon), Three-banded Grasshopper (Banded Range Grasshopper), Differential Grasshopper, Red-legged Locust, Spur-throated Grasshopper, Two-striped Grasshopper, Panther-spotted grasshopper, Southeastern Lubber Grasshopper, Alutacea Bird Grasshopper, American Bird Grasshopper, Green Valley Grasshopper, Toothpick Grasshopper, Horse Lubber Grasshopper, Pallid-winged Grasshopper, and Great Crested Grasshopper (Dinosaur Grasshopper); Long-horned grasshoppers and Katydids family such as Oblong-winged Katydid, Mormon Cricket, Short-legged Shield back Katydid, California Katydid, Angular-winged Katydid, Keeled Shield-back Katydid, Nebraska Cone-head, Gladiator Katydid, Mescalero Shield-back Katydid, True Katydid (Northern Katydid), and Fork-tailed Bush Katydid; Camel Crickets family such as Spotted Camel Cricket, Secret Cove Cricket, Jerusalem Cricket; True Crickets family such as House Cricket (Cricket on the Hearth), Field Cricket, California Tree Cricket, Snowy Tree Cricket, and Black-horned Tree Cricket; Mole Cricket family such as Northern Mole Cricket and European Mole Cricket True Bugs Plant Bug family such as Adelphocoris Plant Bugs, Scarlet Plant bugs, Tarnished Plant Bugs, and Four-lined Plant Bug; Assassin bug family such as Bee Assassins, Wheel Bug; Ambush Bug family such as Jagged Ambush Bug; Lace Bug family such as Oak Lace Bug; Seed Bug family such as Small Eastern Milkweed Bug; Long-necked Seed Bug, and Large Milkweed Bug; Stilt Bug family such as Stilt Bugs; Leaf-footed Bug family such as Florida Leaf-footed Bug, and Squash Bug; Scantless Plant Bug family such as Eastern Boxelder Bug and Western Boxelder Bug; Stink Bug family such as Green Stink Bug, Brochymenas, Two-spotted Stink Bug (Conspicuous Stink Bug), Harlequin Cabbage Bug (Calico Bug, Fire Bug), and Spined Soldier Bug Cicadas and Kin Cicada family such as Periodical Cicadas, Dogday Harvestfly, and Grand Western Cicada Treehoppers Treehopper family, such as Thorn-mimic Treehopper, Oak Treehopper, Buffalo Treehopper, and Locust Treehopper Leafhoppers Leafhopper family such as Grape Leafhopper, Scarlet-and-green		

NOTE: The total number of pages must be divisible by four (4).
The maximum number of pages for this book is forty (40).

BOOKLET PAGES

BOOKLET TEXT (Pages 2-40)

.25"	4.25"	.25"
.25"		
5.5"		6.0"
.25"		
.25"	4.75"	

PESTS CONTROLLED - continued Leafhoppers - continued Leafhopper, Sharpshooter Thrips Banded Thrips family such as Banded-wing Thrips; Common Thrips family Beetles Tiger Beetle family such as Beautiful Tiger Beetle, Dainty Tiger Beetle, Six-Spotted Green Tiger Beetle, and Dejean's Flightless Tiger Beetle; Ground Beetle family such as Bombardier Beetle, Fiery Searcher (Caterpillar hunter), European Ground Beetle, Green Pubescent Ground Beetle, Common Black Ground Beetle, and Boat-backed Ground Beetle; Predacious Diving Beetle family such as Small Flat Diving Beetle, Large Diving Beetle, and Marbled Diving Beetle; Whirligig Beetle family such as Large Whirligig Beetle, and Small Whirligig Beetle; Hister Beetle family such as Hister beetles; Rove Beetle family such as Gold-and-brown Rove Beetle and Pictured Rove Beetle; Carrion Beetle family such as Merined Burying Beetle, Tomentose Burying Beetle (Gold-necked Carrion Beetle), American Carrion Beetle, and Northern Carrion Beetle; Stag Beetle family such as Elephant Stag Beetle, Agassiz's Flat-horned Stag Beetle, Reddish-brown Stag Beetle, and Rugose Stag Beetle; Bessbugs family such as Patent-leather Beetle (Horned Passalus); Scarab Beetle family such as Tumblebugs, Goldsmith Beetle, Green Jumping Beetle, Brown Fruit Chafer, Glossy Pillbug, Grapevine Beetle, May Beetles (June Beetles), Ten-lined June Beetles, and Japanese Beetle; Metallic Wood-Boring Beetles such as Golden Buprestid, Western Pine Borer (Sculptured Pine Borer), and Divergent Metallic Wood Borer (Flatheaded Cherry Tree Borer); Click Beetle family such as Eastern Eyed Click Beetle (Big-eyed Click Beetle), and Fire Beetles; Fireflies (Lightning Bugs) family such as Pyralis Firefly, and Pennsylvania Firefly; Soldier Beetle family such as Pennsylvania Leather-wing, and Downy Leather-wing; Net-winged Beetle family such as Banded Net-wing, and Golden Net-wing; Spider Beetle family such as Spider Beetles, and Texan Spider Beetle; Branch and Twig Borer family such as Apple Twig Borer; Checkered Beetle family such as California Checkered Beetle, Elegant Checkered Beetle, Slender Checkered Beetle, and Red-blue Checkered Beetle; Flat Bark Beetle family such as Red Flat Bark Beetle; Pleasing Fungus Beetle such as Rough Fungus Beetle; Ladybug Beetles such as Two-spotted Ladybug Beetle, Nine-spotted Ladybug Beetle, Spotted "Nine-spotted" Ladybug, Convergent Ladybug Beetle, and Ash Gray Ladybug Beetle; Darkling Beetles such as Horned Fungus Beetle, Broad-necked Darkling Beetle, Plicate Beetle, Yellow Mealworm Beetle, and Ironclad Beetle; Fire-colored Beetles family such as Fire Beetles; Blister Beetles family such as Striped Blister Beetle, Arizona Blister Beetle, Short-winged Blister Beetle (Oil Beetle); Long-horned Beetle family such as Black-horned Pine Borer, Yellow Douglass Fir Borer, Long-jawed Longhorn, Elder Borer (Cloaked Knotty-horn), Twig Pruners, Pine Sawyer (Spined-neck Longhorn), Golden-haired Flower Longhorn, Locust Borer, Black Pine Sawyer (White Spotted Sawyer), Cylindrical Hardwood Borer, Cottonwood Twig Borer, Cottonwood Borer, Giant Root Borers, Ribbed Pine Borer, California Laurel Borer (Banded Alder Borer), Ivory marked Beetle, Red Milkweed Beetle (Eastern Milkweed Longhorn), Notch-tipped Flower Longhorn, and Willow Borer; Seed Borer family such as Bean Weevil; Leaf Beetle family such as Dogwood Calligrapha, Milkweed Tortoise Beetle (Argus Tortoise Beetle), Dogbane Leaf Beetle, Milkweed Leaf Beetle, Willow Leaf Beetle, Spotted Asparagus Beetle, Clavate Tortoise Beetle, Spotted Cucumber Beetle, Waterlily Leaf Beetle, Swamp Milkweed Leaf Beetle, Three-lined Potato Beetle (Old-fashioned Potato Bug), Colorado Potato Beetle, and Girdled Leaf Beetle; Primitive Weevil family such as Oak Timberworm Beetle (Primitive Weevil), Snout Beetles and Weevil family such as Boll Weevil, Black Oak Acorn Weevil, Lesser Cloverleaf Weevil, Alfalfa Weevil, Pine Weevils, Rose Weevils, Agrave Billbug, and Stored-grain Billbug (Granary Weevil, Elephant Bug); Bark and Ambrosia Beetle family such as Red Turpentine Beetle, Pine and Spruce Engraver Beetles, and Bark Beetles	PESTS CONTROLLED - continued Bees, Ants, Wasps, Sawflies Cimbid Sawflies family such as Rusty Willow Sawfly; Common Sawfly family such as Northeastern sawfly; Horntail family such as Pigeon Horntail, and Smokey Horntails; Stem Sawflies family such as Raspberry Horntail; Braconids family such as Braconid Wasps; Ichneumon family such as Giant Ichneumon, Short-tailed Ichneumon, and Red-tailed Ichneumon; Tomyid Wasp family such as California Tomyid; Chalcid family such as Golden-yellow Chalcid; Gall Wasp family such as California Oak Gall Wasps and Live Oak Gall Wasp; Pelecinid family such as American Pellicin; Tiphid Wasp family such as Five-banded Tiphid Wasp; Velvet-Ants such as Thistledown Velvet-ant, Red Velvet-ant, and Cow Killer; Scoliid Wasp family such as Scarab-hunter Wasp, and Digger Wasp; Ant family such as Spine-waisted ants, Leafcutter ants, Texas Carpenter ants, Black Carpenter ants, Crater-nest ants, Crazy Ants, Tawny Crazy Ants, Texas Shed-builder ant (Acrobat ant), Red Ant, Legianary Ant, Little Black Ant, And Lands Honey Ant, Big-headed Ant, Honey Ant, and Fire Ant; Vespid Wasp family such as Potter Wasp, Paper Wasps, Yellow Jackets, and Sandhills Hornet; Spider Wasps such as Blue-black Spider Wasps, and Tarantula Hawk; Sphecoid Wasp family such as Thread-waisted Wasps, Eastern Sand Wasp, Steel-blue Cricket Hunter, Purplish-blue Cricket Hunter, Florida Hunting Wasp, Black and Yellow Mud Dauber, Cicada Killer, and Great Golden Digger Wasp; Bee families such as Plasterer Bee, Yellow-faced Bees, Virescent Green Metallic Bees, Augochora Green Metallic Bees, Alkali Bee, Mining Bee, Willow Mining Bee, Nevada Mining Bee, Clarkia Bee, Faithful Leafcutting Bee, Western Leafcutting Bee, Mason Bees, California Leafcutting Bee, Digger Bees, Western Cuckoo Bee, California Carpenter Bee Flies Crane fly family such as Wood-boring Tupidid, Giant Western Crane Fly, and Crane Flies; Phantom Crane Flies such as Phantom Crane Fly; Net-winged Midges such as Comstock's Net-winged Midge; Phantom Midges; Black Flies such as Black flies; March Flies family such as March Flies; Fungus Gnats family such as Fungus Gnats; Snipe Fly family such as Gold-backed Snipe Fly; Stiletto Fly family such as Stiletto Flies; Flower-loving Fly family such as Flower-loving Fly; Mydas Fly family such as Mydas fly; Robber Fly family such as Bearded Robber Fly, Sacken's Bee Hunter, Giant Robber Fly, Bee Killer (Giant Robber Fly), and Robber Flies; Bee Flies such as Bee Fly, Large Bee Flies, and Progressive bee Flies; Dance Flies such as Dance Flies; Long-Legged Flies such as Condylostylid Long-legged Flies, and Texan Long-legged Fly; Hover Fly family such as Woolly Bear Hover Flies, Elongate Aphid Flies, Drone Flies, Bulb Flies, American Hover Flies, and Toxomerus Hover Fly; Thick Headed Fly family such as Thick headed fly; Pyrgoid Fly family such as Pyrgoid Flies; Fruit Fly family such as Walnut Husk Fly, and Apple Maggot Fly; Seaweed Fly family such as California Seaweed Fly; Marsh Fly family such as Marsh flies; Shore fly family such as Yellowstone Brine Fly; Pomace fly family such as Vinegar Fly; Anthomyid Fly family such as Dung Fly; Muscid Fly family such as House Fly, and Biting Stable Fly; Louse Fly family, Blow Fly family such as Blue Bottle Fly, Screw-worm Fly, and Green Bottle Fly; Flesh Fly family such as Flesh flies; Tachinid Fly family such as Beelike Tachinid Fly, Early Tachinid Fly, Repetitive Tachinid Fly, Tachina Fly, and Fringe-legged Tachinid Fly Spiders Folding trapdoor spider family such as Turret spider; Tarantula family such as Desert Tarantula; Trapdoor Spider family such as California Trapdoor spider; Dichnid spider family such as Branch-ypid spiders; Spitting spider family such as Spitting spiders; Violin Spider family such as Desert Loxosceles, and Violin Spider; Comb-footed Spider family such as American House Spider, and Black Widow Spider, Sheet-web Weaver family such as Hammock Spider; Orb Weaver family such as Orb weavers, barn spider, Garden spider (cross spider), Marbled orb Weaver, Shamrock Spider, Six-spotted orb-weaver, Silver Argiope, Black-and-Yellow Argiope, Crablike Spiny Orb Weaver, Bola Spider, Arrow-shaped Micrathena, and Golden-silk spider (calico spider);
--	--

BOOKLET PAGES

BOOKLET TEXT (Pages 2-40)

.25"		4.25"		.25"																																																																																			
.25"		5.5"		6.0"																																																																																			
.25"																																																																																							
<table><tr><th colspan="3">PESTS CONTROLLED - continued</th><th>Plants</th><th>Pests</th><th>Use Before Harvest</th></tr><tr><td colspan="3">Spiders - continued</td><td>VEGETABLES</td><td></td><td></td></tr><tr><td colspan="3">Large-jawed Orb Weaver family such as Mabel Orchard Spider, Venusta Orchard Spider, Elongate Long-jawed Orb Weaver, and Long-jawed Orb Weaver; Funnel Web Weaver family such as Grass Spiders; Nursery Web Spider family such as Brownish-grey Fishing Spider, Six-spotted Fishing spider, and Nursery Web Spider; Wolf Spider family such as Burrowing Wolf Spiders, Carolina Wolf spider, Forest Wolf spider, rabid wolf spider, and Thin-legged Wolf spider; Lynx Spider family such as Jumping Lynx spider, and green lynx spider; Sac Spider family such as Ant-mimic spiders; Wandering spider family such as Wandering spiders; Giant Crab spider family such as Huntsman spider, and Golden Huntsman spider; Selenopid Crab Spider family such as Selenopid Crab Spider; Crab Spider family such as Goldenrod spider (flower spider, red-spotted Crab Spider), Elegant Crab Spider, Thrice-banded Crab Spider; Philodromid family such as inconspicuous Crab Spiders; Jumping Spiders such as Green Lyssomanes, Dimorphic Jumping Spider, Metaphid Jumping Spider, Ant-mimic Jumping spider, and Daring Jumping spider</td><td> • Groundcherry (<i>Physalis</i> spp.); • Pepino (Melon pear); • Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper); • Tomatillo; • Tomato </td><td>Colorado Potato Beetle, Corn Borer, European, Corn Borer, Southwestern Corn Earworm, Cucumber Beetle, Cutworm, Flea Beetle, Garden Webworm, Green Stink Bug, Hornworms, Leafminers (adults), Leafhoppers, Meadow Spittlebug, Pepper Maggot (adults), Pepper Weevil, Plant Bugs, Tobacco Budworm, Tomato Fruitworm, Tomato Pinworm, Aphids, Armyworm, Beet Armyworm, Fall Cabbage Looper, Grasshoppers, Lygus Bugs, Brown Stink Bug, Tomato Psyllid, Thrips, Whiteflies</td><td></td></tr><tr><td colspan="3">Daddy-long-legs family such as Eastern Daddy-long-legs and Brown Daddy-long-legs</td><td>Brassica Vegetables including:</td><td>Corn Earworm, Cucumber Beetles, Cutworm, Diamondback Moth, Flea Beetles, Imported Cabbageworm, Leafhoppers, Saltmarsh Caterpillar, Southern Cabbageworm, Tobacco Budworm, Alfalfa Looper, Armyworms, Cabbage Looper, Cabbage Webworm, Ground Beetles, Leafminers (adults), Lygus Bugs, Onion Thrips, Stinkbugs, Wireworm (adults), Aphids, and Whiteflies</td><td rowspan="2">Spray up to 1 day before harvesting</td></tr><tr><td colspan="3">Ticks</td><td> • Broccoli; • Chinese Broccoli (gai lan, white flowering broccoli); • Brussels Sprouts; • Cauliflower; • Cavalo Broccolo; • Kohlrabi; • Cabbage; • Chinese Cabbage (napa); • Leafy Brassica Greens including: Broccoli Raab (rapini); • Chinese Cabbage (bok choy); • Chinese Mustard Cabbage (gai choy); • Collards; • Kale; • Mizuna; • Mustard Greens; • Mustard Spinach; • Rape Greens </td><td></td></tr><tr><td colspan="3">Psyllid, Whiteflies, Aphids, Mealybugs, Scale Insects</td><td>Root & Tuber Vegetables</td><td>Cutworms, Cabbage Looper, Cucumber Beetle, European Corn Borer, Flea beetles, Leafhoppers, Southern Corn Rootworm (adult), Vegetable Weevil, Whitefringed Beetle (adult), Armyworm, Beet Armyworm, Yellowstriped Cabbage Maggot, Colorado Potato Beetle, Grasshoppers, Imported Cabbageworm, Potato Leafhopper, Potato Psyllid, Potato Tuberworm, Tarnished Plant Bug</td><td rowspan="2">Spray up to 1 day before harvesting</td></tr><tr><td colspan="3">Psyllid family such as American Alder Psyllid; Whitefly family such as Greenhouse Whitefly; Aphid family such as Green Apple Aphid, Rosy Apple Aphid, Rose, Pea and Potato Aphid, Root Aphid, Coudy-winged Cottonwood Aphid, and Giant Willow Aphid; Woolly and Gail-Making Aphid family such as Woolly Apple Aphid; Giant Scale Insects family such as Cottony Cushion Scale; Armored Scale Insects family such as Oyster Shell Scale; Mealybug family such as Long-tailed mealybug; Cochineal Bug family such as Cochineal bug</td><td> • Arracacha; • Arrowroot; • Artichoke; • Garden Beet; • Edible Burdock; • Edible Canna; • Carrot; • Turnip-rooted Chervil; • Chicory; • Chufa; • Dasheen (Taro); • Ginger; • Horseradish; • Leren; </td><td></td></tr><tr><td colspan="3">Millipedes</td><td>Fruiting Vegetables including:</td><td>Armyworm, Southern Armyworm, True Armyworm, Yellow-striped Celery Leaf Tier,</td><td>Spray up to 1 day before harvesting</td></tr><tr><td colspan="3">Millipedes such as soft millipedes, Florida millipede, Almond millipede, North American millipede and flat millipedes</td><td></td><td></td><td></td></tr><tr><td colspan="3"></td><td>Plants</td><td>Pests</td><td>Use Before Harvest</td></tr><tr><td colspan="3">VEGETABLES</td><td>Leafy Vegetables</td><td>Com Earworm Cucumber Beetles, Cutworms, Diamondback Moth, Flea Beetles Imported Cabbageworm, Leafhoppers, Saltmarsh Caterpillar, Tobacco Budworm, Armyworms, Loopers, Lygus Bugs, Onion Thrips, Stink Bugs, Wireworm</td><td>Spray up to 1 day before harvesting</td></tr><tr><td colspan="3"> • Amaranth (leafy amaranth, Chinese spinach, tampala); • Arugula; • Cardoon; • Cilantro • Corn Salad; • Garden Cress; • Upland Cress; • Dandelion; • Dock; • Endive; • Florence Fennel; • Head Lettuce; • Leaf Lettuce; • Orach; • Parsley; • Garden Purslane; • Winter Purslane; • Radicchio (red chicory); • Rhubarb; • Spinach; • Swiss Chard </td><td></td><td></td><td></td></tr><tr><td colspan="3">Fruiting Vegetables including:</td><td>Armyworm, Southern Armyworm, True Armyworm, Yellow-striped Celery Leaf Tier,</td><td></td><td></td></tr><tr><td colspan="3">• Eggplant;</td><td></td><td></td><td></td></tr></table>						PESTS CONTROLLED - continued			Plants	Pests	Use Before Harvest	Spiders - continued			VEGETABLES			Large-jawed Orb Weaver family such as Mabel Orchard Spider, Venusta Orchard Spider, Elongate Long-jawed Orb Weaver, and Long-jawed Orb Weaver; Funnel Web Weaver family such as Grass Spiders; Nursery Web Spider family such as Brownish-grey Fishing Spider, Six-spotted Fishing spider, and Nursery Web Spider; Wolf Spider family such as Burrowing Wolf Spiders, Carolina Wolf spider, Forest Wolf spider, rabid wolf spider, and Thin-legged Wolf spider; Lynx Spider family such as Jumping Lynx spider, and green lynx spider; Sac Spider family such as Ant-mimic spiders; Wandering spider family such as Wandering spiders; Giant Crab spider family such as Huntsman spider, and Golden Huntsman spider; Selenopid Crab Spider family such as Selenopid Crab Spider; Crab Spider family such as Goldenrod spider (flower spider, red-spotted Crab Spider), Elegant Crab Spider, Thrice-banded Crab Spider; Philodromid family such as inconspicuous Crab Spiders; Jumping Spiders such as Green Lyssomanes, Dimorphic Jumping Spider, Metaphid Jumping Spider, Ant-mimic Jumping spider, and Daring Jumping spider			• Groundcherry (<i>Physalis</i> spp.); • Pepino (Melon pear); • Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper); • Tomatillo; • Tomato	Colorado Potato Beetle, Corn Borer, European, Corn Borer, Southwestern Corn Earworm, Cucumber Beetle, Cutworm, Flea Beetle, Garden Webworm, Green Stink Bug, Hornworms, Leafminers (adults), Leafhoppers, Meadow Spittlebug, Pepper Maggot (adults), Pepper Weevil, Plant Bugs, Tobacco Budworm, Tomato Fruitworm, Tomato Pinworm, Aphids, Armyworm, Beet Armyworm, Fall Cabbage Looper, Grasshoppers, Lygus Bugs, Brown Stink Bug, Tomato Psyllid, Thrips, Whiteflies		Daddy-long-legs family such as Eastern Daddy-long-legs and Brown Daddy-long-legs			Brassica Vegetables including:	Corn Earworm, Cucumber Beetles, Cutworm, Diamondback Moth, Flea Beetles, Imported Cabbageworm, Leafhoppers, Saltmarsh Caterpillar, Southern Cabbageworm, Tobacco Budworm, Alfalfa Looper, Armyworms, Cabbage Looper, Cabbage Webworm, Ground Beetles, Leafminers (adults), Lygus Bugs, Onion Thrips, Stinkbugs, Wireworm (adults), Aphids, and Whiteflies	Spray up to 1 day before harvesting	Ticks			• Broccoli; • Chinese Broccoli (gai lan, white flowering broccoli); • Brussels Sprouts; • Cauliflower; • Cavalo Broccolo; • Kohlrabi; • Cabbage; • Chinese Cabbage (napa); • Leafy Brassica Greens including: Broccoli Raab (rapini); • Chinese Cabbage (bok choy); • Chinese Mustard Cabbage (gai choy); • Collards; • Kale; • Mizuna; • Mustard Greens; • Mustard Spinach; • Rape Greens		Psyllid, Whiteflies, Aphids, Mealybugs, Scale Insects			Root & Tuber Vegetables	Cutworms, Cabbage Looper, Cucumber Beetle, European Corn Borer, Flea beetles, Leafhoppers, Southern Corn Rootworm (adult), Vegetable Weevil, Whitefringed Beetle (adult), Armyworm, Beet Armyworm, Yellowstriped Cabbage Maggot, Colorado Potato Beetle, Grasshoppers, Imported Cabbageworm, Potato Leafhopper, Potato Psyllid, Potato Tuberworm, Tarnished Plant Bug	Spray up to 1 day before harvesting	Psyllid family such as American Alder Psyllid; Whitefly family such as Greenhouse Whitefly; Aphid family such as Green Apple Aphid, Rosy Apple Aphid, Rose, Pea and Potato Aphid, Root Aphid, Coudy-winged Cottonwood Aphid, and Giant Willow Aphid; Woolly and Gail-Making Aphid family such as Woolly Apple Aphid; Giant Scale Insects family such as Cottony Cushion Scale; Armored Scale Insects family such as Oyster Shell Scale; Mealybug family such as Long-tailed mealybug; Cochineal Bug family such as Cochineal bug			• Arracacha; • Arrowroot; • Artichoke; • Garden Beet; • Edible Burdock; • Edible Canna; • Carrot; • Turnip-rooted Chervil; • Chicory; • Chufa; • Dasheen (Taro); • Ginger; • Horseradish; • Leren;		Millipedes			Fruiting Vegetables including:	Armyworm, Southern Armyworm, True Armyworm, Yellow-striped Celery Leaf Tier,	Spray up to 1 day before harvesting	Millipedes such as soft millipedes, Florida millipede, Almond millipede, North American millipede and flat millipedes									Plants	Pests	Use Before Harvest	VEGETABLES			Leafy Vegetables	Com Earworm Cucumber Beetles, Cutworms, Diamondback Moth, Flea Beetles Imported Cabbageworm, Leafhoppers, Saltmarsh Caterpillar, Tobacco Budworm, Armyworms, Loopers, Lygus Bugs, Onion Thrips, Stink Bugs, Wireworm	Spray up to 1 day before harvesting	• Amaranth (leafy amaranth, Chinese spinach, tampala); • Arugula; • Cardoon; • Cilantro • Corn Salad; • Garden Cress; • Upland Cress; • Dandelion; • Dock; • Endive; • Florence Fennel; • Head Lettuce; • Leaf Lettuce; • Orach; • Parsley; • Garden Purslane; • Winter Purslane; • Radicchio (red chicory); • Rhubarb; • Spinach; • Swiss Chard						Fruiting Vegetables including:			Armyworm, Southern Armyworm, True Armyworm, Yellow-striped Celery Leaf Tier,			• Eggplant;					
PESTS CONTROLLED - continued			Plants	Pests	Use Before Harvest																																																																																		
Spiders - continued			VEGETABLES																																																																																				
Large-jawed Orb Weaver family such as Mabel Orchard Spider, Venusta Orchard Spider, Elongate Long-jawed Orb Weaver, and Long-jawed Orb Weaver; Funnel Web Weaver family such as Grass Spiders; Nursery Web Spider family such as Brownish-grey Fishing Spider, Six-spotted Fishing spider, and Nursery Web Spider; Wolf Spider family such as Burrowing Wolf Spiders, Carolina Wolf spider, Forest Wolf spider, rabid wolf spider, and Thin-legged Wolf spider; Lynx Spider family such as Jumping Lynx spider, and green lynx spider; Sac Spider family such as Ant-mimic spiders; Wandering spider family such as Wandering spiders; Giant Crab spider family such as Huntsman spider, and Golden Huntsman spider; Selenopid Crab Spider family such as Selenopid Crab Spider; Crab Spider family such as Goldenrod spider (flower spider, red-spotted Crab Spider), Elegant Crab Spider, Thrice-banded Crab Spider; Philodromid family such as inconspicuous Crab Spiders; Jumping Spiders such as Green Lyssomanes, Dimorphic Jumping Spider, Metaphid Jumping Spider, Ant-mimic Jumping spider, and Daring Jumping spider			• Groundcherry (<i>Physalis</i> spp.); • Pepino (Melon pear); • Pepper (includes bell pepper, chili pepper, cooking pepper, pimento, sweet pepper); • Tomatillo; • Tomato	Colorado Potato Beetle, Corn Borer, European, Corn Borer, Southwestern Corn Earworm, Cucumber Beetle, Cutworm, Flea Beetle, Garden Webworm, Green Stink Bug, Hornworms, Leafminers (adults), Leafhoppers, Meadow Spittlebug, Pepper Maggot (adults), Pepper Weevil, Plant Bugs, Tobacco Budworm, Tomato Fruitworm, Tomato Pinworm, Aphids, Armyworm, Beet Armyworm, Fall Cabbage Looper, Grasshoppers, Lygus Bugs, Brown Stink Bug, Tomato Psyllid, Thrips, Whiteflies																																																																																			
Daddy-long-legs family such as Eastern Daddy-long-legs and Brown Daddy-long-legs			Brassica Vegetables including:	Corn Earworm, Cucumber Beetles, Cutworm, Diamondback Moth, Flea Beetles, Imported Cabbageworm, Leafhoppers, Saltmarsh Caterpillar, Southern Cabbageworm, Tobacco Budworm, Alfalfa Looper, Armyworms, Cabbage Looper, Cabbage Webworm, Ground Beetles, Leafminers (adults), Lygus Bugs, Onion Thrips, Stinkbugs, Wireworm (adults), Aphids, and Whiteflies	Spray up to 1 day before harvesting																																																																																		
Ticks			• Broccoli; • Chinese Broccoli (gai lan, white flowering broccoli); • Brussels Sprouts; • Cauliflower; • Cavalo Broccolo; • Kohlrabi; • Cabbage; • Chinese Cabbage (napa); • Leafy Brassica Greens including: Broccoli Raab (rapini); • Chinese Cabbage (bok choy); • Chinese Mustard Cabbage (gai choy); • Collards; • Kale; • Mizuna; • Mustard Greens; • Mustard Spinach; • Rape Greens																																																																																				
Psyllid, Whiteflies, Aphids, Mealybugs, Scale Insects			Root & Tuber Vegetables	Cutworms, Cabbage Looper, Cucumber Beetle, European Corn Borer, Flea beetles, Leafhoppers, Southern Corn Rootworm (adult), Vegetable Weevil, Whitefringed Beetle (adult), Armyworm, Beet Armyworm, Yellowstriped Cabbage Maggot, Colorado Potato Beetle, Grasshoppers, Imported Cabbageworm, Potato Leafhopper, Potato Psyllid, Potato Tuberworm, Tarnished Plant Bug	Spray up to 1 day before harvesting																																																																																		
Psyllid family such as American Alder Psyllid; Whitefly family such as Greenhouse Whitefly; Aphid family such as Green Apple Aphid, Rosy Apple Aphid, Rose, Pea and Potato Aphid, Root Aphid, Coudy-winged Cottonwood Aphid, and Giant Willow Aphid; Woolly and Gail-Making Aphid family such as Woolly Apple Aphid; Giant Scale Insects family such as Cottony Cushion Scale; Armored Scale Insects family such as Oyster Shell Scale; Mealybug family such as Long-tailed mealybug; Cochineal Bug family such as Cochineal bug			• Arracacha; • Arrowroot; • Artichoke; • Garden Beet; • Edible Burdock; • Edible Canna; • Carrot; • Turnip-rooted Chervil; • Chicory; • Chufa; • Dasheen (Taro); • Ginger; • Horseradish; • Leren;																																																																																				
Millipedes			Fruiting Vegetables including:	Armyworm, Southern Armyworm, True Armyworm, Yellow-striped Celery Leaf Tier,	Spray up to 1 day before harvesting																																																																																		
Millipedes such as soft millipedes, Florida millipede, Almond millipede, North American millipede and flat millipedes																																																																																							
			Plants	Pests	Use Before Harvest																																																																																		
VEGETABLES			Leafy Vegetables	Com Earworm Cucumber Beetles, Cutworms, Diamondback Moth, Flea Beetles Imported Cabbageworm, Leafhoppers, Saltmarsh Caterpillar, Tobacco Budworm, Armyworms, Loopers, Lygus Bugs, Onion Thrips, Stink Bugs, Wireworm	Spray up to 1 day before harvesting																																																																																		
• Amaranth (leafy amaranth, Chinese spinach, tampala); • Arugula; • Cardoon; • Cilantro • Corn Salad; • Garden Cress; • Upland Cress; • Dandelion; • Dock; • Endive; • Florence Fennel; • Head Lettuce; • Leaf Lettuce; • Orach; • Parsley; • Garden Purslane; • Winter Purslane; • Radicchio (red chicory); • Rhubarb; • Spinach; • Swiss Chard																																																																																							
Fruiting Vegetables including:			Armyworm, Southern Armyworm, True Armyworm, Yellow-striped Celery Leaf Tier,																																																																																				
• Eggplant;																																																																																							
.25"		4.75"																																																																																					

NOTE: The total number of pages must be divisible by four (4).

The maximum number of pages for this book is forty (40).

BOOKLET TEXT
(Pages 2-40)

NOTE: The total number of pages must be divisible by four (4).
The maximum number of pages for this book is forty (40).

NOTE: The total number of pages must be divisible by four (4).
The maximum number of pages for this book is forty (40).

BOOKLET PAGES

BOOKLET TEXT (Pages 2-40)

.25"		4.25"		.25"									
.25"		PRIMEROS AUXILIOS En caso de ingestión • Llame inmediatamente a un centro de control de envenenamiento o un médico para recibir consejos sobre el tratamiento. • Si la persona puede tragar, haga que beba un vaso de agua. • No induzca el vómito a menos que lo recomiende el centro de control de envenenamiento o el médico. • No le de nada por la boca a una persona inconsciente. En caso de contacto con los ojos • Mantenga el ojo abierto y enjuáguelo lenta y suavemente con agua durante 15-20 minutos. • Si usa lentes de contacto, retírelos después de los primeros 15 minutos y luego continúe enjuagando el ojo. • Llame a un centro de control de envenenamiento o un médico para recibir consejos sobre el tratamiento. TELÉFONO DE ASISTENCIA DE EMERGENCIA Tenga a mano el recipiente o la etiqueta del producto cuando llame al centro de control de envenenamiento o al médico, o cuando acuda para recibir tratamiento. También puede llamar al servicio de asistencia de emergencia al 1-800-420-9347. NOTA PARA EL MÉDICO El contacto con la piel (vapor o polvo) puede anteceder a una sensación de cosquilleo o entumecimiento, usualmente de la cara, pero pasa sin necesidad de tratamiento. Este producto es un piretroides. En caso de ingestión de una gran cantidad, el estómago y los intestinos deben ser evacuados. El tratamiento es sintomático y paliativo. Para más información sobre el uso de este producto, llame al 1-866-945-5033. PRECAUCIONES PELIGROS AMBIENTALES Este pesticida es extremadamente tóxico para peces, invertebrados acuáticos, ostras y camarones. Para proteger el ambiente, no permita que el pesticida entre o sea arrastrado hacia desagües para tormentas, zanjas de drenaje, canaletas ni aguas superficiales. Aplique el producto cuando haga buen tiempo y no se pronostique lluvia en las próximas 24 horas, esto evitará que el viento o la lluvia vuelen o arrastren el pesticida hacia afuera del área tratada. Enjuague el equipo de aplicación sobre el área tratada para evitar que el agua usada llegue a cuerpos de agua o sistemas de drenajes. Se debe tener cuidado al rociar para evitar que afecte a peces y reptiles domésticos de los estanques ornamentales. Este producto es altamente tóxico para las abejas expuestas al tratamiento directo o los residuos en malezas o cultivos floridos. No aplique este producto ni permita que caiga sobre malezas o plantas floridas si hay abejas forrajeando en el área de tratamiento. INDICACIONES DE USO El uso de este producto de cualquier manera que difiera de lo indicado en su etiqueta constituye una violación de las leyes federales. AGÍTELO BIEN ANTES DE USARLO RESTRICCIONES DE USO No use este producto para el tratamiento de mascotas. No aplique este producto de manera que haga contacto con adultos, niños o mascotas, ya sea directa o accidentalmente. No permita que los adultos, niños ni mascotas entren al área tratada hasta que se seque el rociado. Retire las mascotas y aves, y cubra los acuarios de peces antes de rociar el producto. Evite la contaminación de los alimentos y piensos. Retire o cubra los alimentos y el agua potable expuestos antes de la aplicación. No lo use en establecimientos donde se manipulen, procesen, preparen y/o sirvan alimentos. Cuando lo use en el hogar, retire o cubra los platos, utensilios, equipos para procesar alimentos y superficies para preparar alimentos, o lávelos antes de utilizarlos. No lo aplique directamente a los animales. No lo use en el interior de los aviones. PARA USOS EN EXTERIORES Todas las aplicaciones en exteriores deben limitarse al tratamiento localizado o de lugares angostos solamente, excepto para los siguientes usos permitidos: (1) Tratamiento en la tierra o la vegetación alrededor de las estructuras (2) Aplicaciones en céspedes y tepes o en otra vegetación (3) Aplicaciones en los cimientos de las edificaciones hasta una altura máxima de 3 pies (0.91 m). Exceptuando las aplicaciones en los cimientos de las edificaciones, todas las aplicaciones exteriores en superficies impermeables como las aceras, entradas para carros, patios, porches y superficies estructurales (ventanas, puertas y aleros) se limitan al tratamiento localizado y en lugares angostos solamente. No riegue el área tratada hasta el punto en que el producto pueda ser arrastrado por el agua. No lo aplique durante la lluvia. Se prohíbe aplicarlo directamente en alcantarillados o desagües, o cualquier otro lugar, como las canaletas, que puedan drenar el agua hacia alcantarillados, desagües para tormentas, cuerpos de agua o hábitats acuáticos. No permita que este producto entre en ningún desagüe durante o después de la aplicación. Si tiene una situación de emergencia médica, necesita consultar sobre un diagnóstico, terapia o análisis, llame por favor al 1-(800) 420-9347. Si tiene preguntas acerca de la aplicación o el uso del producto, llame por favor al 1-(866) 945-5033. Si se produce un derrame o situación de emergencia durante la transportación, llame por favor al 1-(800) 424-9300. Para los reembolsos a consumidores: TechPac, LLC PO Box 95437 Palatine, IL 60095-0437 <tr><td colspan="2">5.5"</td><td colspan="2">6.0"</td></tr> <tr><td colspan="2">.25"</td><td colspan="2">4.75"</td><td colspan="2">.25"</td></tr>		5.5"		6.0"		.25"		4.75"		.25"	
5.5"				6.0"									
.25"		4.75"		.25"									

NOTE: The total number of pages must be divisible by four (4).
The maximum number of pages for this book is forty (40).

BOOKLET PAGES

BOOKLET TEXT (Pages 2-40)

.25"		4.25"		.25"		1 día (jugosas)													
.25"		5.5"		6.0"		10"													
INSTRUCCIONES PARA MEDIR		<table><tr><th>Onzas líquidas del concentrado producto</th><th>Cantidad de agua</th></tr><tr><td>1</td><td>1 cuarto (32 onzas)</td></tr><tr><td>2</td><td>½ galón (64 onzas)</td></tr><tr><td>4</td><td>1 galón (128 onzas)</td></tr><tr><td>6</td><td>1 ½ galones (192 onzas)</td></tr><tr><td>8</td><td>2 galones (256 onzas)</td></tr></table>		Onzas líquidas del concentrado producto	Cantidad de agua	1	1 cuarto (32 onzas)	2	½ galón (64 onzas)	4	1 galón (128 onzas)	6	1 ½ galones (192 onzas)	8	2 galones (256 onzas)	VEGETALES		PHI	
Onzas líquidas del concentrado producto	Cantidad de agua																		
1	1 cuarto (32 onzas)																		
2	½ galón (64 onzas)																		
4	1 galón (128 onzas)																		
6	1 ½ galones (192 onzas)																		
8	2 galones (256 onzas)																		
 ROCIADORES DE TANQUE TIPO BOMBA		- Cierre firmemente el rociador y genere presión con la bomba de mano. Para las aplicaciones localizadas, use una boquilla de chorro grueso. Para aplicaciones en áreas más grandes, use una boquilla de rociado en abanico. - Sostenga la punta de la vara a 12 pulgadas (30,48 cm) del área de tratamiento mientras rocía.		Legumbres — guisantes de vainas comestibles jugosos, guisantes sin cáscara jugosos y guisantes sin cáscara secos: Guisante enano, guisante de vaina comestible, guisante mollar, tirabeque dulce, gandul, petipúas, arveja, chícharo verde y lenteja		1 día (jugosas)													
				Legumbres — frijoles de vainas comestibles jugosos, frijoles sin cáscara jugosos y frijoles sin cáscara secos: Ayocote, habichuela verde, alubia, frijol espárrago, frijol largo chino, Moth Bean (Vigna aconitifolia), frijol serpiente, frijol machete, frijol de soja (semilla inmadura), haba espada, haba de Lima, haba, frijol de carita, carilla, lupino, lupino dulce, lupino blanco, lupino blanco dulce, haba, frijol colorado, judía, frijol pinto, frijol tépari, frijol adzuki, Catjang (Vigna unguiculata cylindrica), frijol Castilla, alubia Crowder, Moth Bean (Vigna aconitifolia), judía mungo o frijol chino, Rice Bean (Vigna umbellata), frijol negro o lenteja negra, garbanzo, guar, frijol de Egipto		1 día (jugosas)													
 ROCIADORES ESTILO BOTÓN PARA ACOPLAR A LA MANGUERA		- Rellene el frasco del rociador con suficiente producto para el área a tratar. NO AGREGUE AGUA. - Cierre el rociador, regule el botón giratorio, acóplelo a la manguera del jardín y aplíquelo. - Vierta el producto sin usar en el recipiente original del Sevin® Insect Killer Concentrate.		Vegetales que dan frutos: Berenjena, uchuva (Physalis), oca, pepino melón (pepino dulce o melón de árbol), pimiento (incluyendo el pimiento o ají, pimiento chile, pimiento de cocinar, pimiento y pimiento dulce), tomate y tomate		1 día													
				Vegetales de hojas comestibles: Amaranto (amaranto de hojas frondosas, espinaca china y amaranto tricolor); arugula (Rouquette), cardo, apio, apio chino, lechuga china o apio lechuga, perifollo, crisantemo (de hojas comestibles y de guirnalda), cilantro (no se debe usar en cilantro cultivado para semillas o condimento), canónigo, berro (berro o mastuerzo, berro de montaña (barbárea amarilla, berro de invierno)), diente de león, romaza (vinagrera), endivia (endivia de hoja ancha), hinojo de Florencia, berro o mastuerzo, lechuga (de cogollo o beluga y lechuga de hojas), abanico, perejil, verdolaga (de jardín), verdolaga (de invierno), radichio (achicoria roja), ruibarbo, espinaca (incluyendo la de Nueva Zelanda y la enredadera, espinaca de Malabar y la espinaca de la India) y acelga		1 día													
INSTRUCCIONES PARA LA APLICACIÓN		- Aplíquelo donde se espere que aparezcan los insectos o cuando se observen los insectos o el daño. Repita la aplicación si es necesario según las indicaciones específicas del lugar mencionadas abajo. - Aplíquelo en tiempo de calma. - Comience a rociar desde el punto más lejano del área de tratamiento y trabaje hacia atrás para evitar el contacto con las superficies recién tratadas. - Rocielo abundantemente para mojar las superficies de la planta hasta el punto en que el líquido corra. - Las personas o mascotas no deben entrar al área tratada hasta que se seque el rociado. - No lo aplique en árboles de más de 10 pies (3,04 m) de altura. Para ello, considere contratar a un profesional certificado.		Manises		7 días													
		Tubérculos y vegetales de raíces (excepto la remolacha de azúcar): Arracacha, maranta, alcachofa (china y tupinambo), remolacha de huerto, lampazo, platanillo comestible, zanahoria, yuca (amarga y dulce), apio (raíz), chayote (raíz), vegetal de la raíz del perifollo, achicoria, chufa, malanga (Taro), jengibre, ginseng, rábano picante, lerén, vegetal de la raíz del perejil, chirivía, papa, rábano blanco o rábano japonés (daikon), rábano, colinabo, salsifi (barba cabruna), escorzonera negra, escorzonera, escaravía, camote, yautía, cúrcuma, nabo, jicama, ñame		1 día															
		Girasoles: Ricino o higuera, árbol de sebo, Euphorbia, onagra, jojoba, negrillo, escaramujo o rosa mosqueta, áster Stokesia, eucalipto de sebo, árbol del aceite de té o de camelia y áster Vernonia		30 días															
		BAYAS, FRUTAS Y NUECES																	
		Use el Sevin® Concentrate Insect Killer como tratamiento localizado para las infestaciones de plagas en las plantas de bayas, frutas y nueces mencionadas abajo.																	
		No se recomienda la aplicación en árboles de más de 10 pies (3,04 m) ni árboles con áreas inaccesibles para ser rociadas.																	
		Aplique la mezcla de rociado a las hojas superiores e inferiores, alrededor de los ramilletes de frutos y en las ramas y troncos hasta que el líquido corra. Repita la aplicación si regresan los insectos, pero no más de una vez cada siete días y según el intervalo previo a la cosecha.																	
				10"															
.25"		4.75"		.25"															

**NOTE: The total number of pages must be divisible by four (4).
The maximum number of pages for this book is forty (40).**

BOOKLET TEXT

(Pages 2-40)

BAYAS, FRUTAS Y NUECES	PHI	ORUGAS, MARIPOSAS Y POLILLAS
Grupo de las bayas incluyendo: Mora, mora roja, framuesa roja y negra, arándano azul (de arbusto alto y de arbusto bajo), grosella, baya del saúco, uva crespita o grosella espinosa, arándano, variedades y/o híbridos de estas	1 día	Ermine como la polilla del gusano tejedor <i>Aliaxius</i> ; la familia de polillas de alas transparentes como el bórbor del marah, el bórbor de la enredadera de calabaza, bórbor del álamo, bórbor del sicomoro de California; polillas de la familia de los tortricidos como el gusano enrollador de hojas de frutales, gusano tóxico del naranjo y polilla del manzano; hespéridos como la mariposa de manchas plateadas y mariposa de cola larga; mariposas de cola de galandina como la mariposa verde, mariposa cebrina, cola de galandina gigante, mariposa tigre, cola de galandina negra del este, <i>Speiobus swallowtail</i> (<i>Papilio troilus</i>); y <i>Phoebus</i> ; mariposas blancas y anaranjadas de la familia antiochani como la antiochani sara, antiochani grande del sur y la mariposa del repollo europeo; mariposas de la familia leucénidos como la mariposa morada, Spring Azure (<i>Calastina ladea</i>), mariposa de color azul del este, cosechadora, mariposa azul plateada, mariposa Brown Elfín, azul marino, <i>Lycaena phlaeas</i> y mariposa grise; mariposas de la familia Riodinidae como la mariposa del norte; mariposas de la familia de libéllos como la mariposa común; mariposas de la familia de ninfalidos como la fritillaria del golfo, white peacock (<i>Anartia jatrophae</i>); Harris Checkerspot (<i>Chrysobothris harrisii</i>), dama pintada, cebrina de Baltimore, mariposa virrey, admiral blanca, Mourning Cloak (<i>Nymphalis antiopa</i>), Milbert's Tortoise Shell (<i>Aglais milberti</i>), mariposa perlada, Question Mark (<i>Polytonia interrogationis</i>), Buckeye (<i>Junonia coenia</i>), Great Spangled Frythia (<i>Speyeria cybele</i>) y admiral roja; mariposas de la familia Asterocampa y <i>Anaeni</i> como la mariposa Hackberry; familia Satyrs, ninfas y ártica como la Wood Nymph (<i>Cercyonis pegala</i>), American Grayling, la Satyr blanca, mariposa marón común, mariposa de ojo perlado y la <i>Chryxus</i> ; mariposas de la familia Danaidae como la reina y la monarca; polillas de la familia de los zigénidos como la polilla de la hoja de la vid; polillas de la familia de los limacodidos como la polilla de la ruga <i>Saddleback</i> ; familia de los pírididos como la polilla mediterránea, polilla tejedora de tierra, polilla Indian Meal, polilla meal, familia <i>Pterophoridae</i> como la Arctioche Plume Mot (<i>Platyptilia carduactylus</i>); polillas de la familia de los gusanos medidores como la polilla Tulip Tree Beauty, polilla del pulgón de California y polilla del gusano medidor de California; polillas de la familia de orugas que forman tiendas como la polilla de la oruga de tienda del oeste; familia de la polilla gigante de la seda como la polilla Luna, polilla Polyphemus, polilla lo, polilla Promethea, polilla real (polilla real del nogal), polilla rosada del arce, polilla imperial, polilla de seda común, polilla Cecropia y polilla <i>Cynthia</i> ; familia de polillas esfíngidos como la esfíngide trepadora de Virginia, esfíngide Pandora, familia de las alas transparentes, familia de las líneas blancas, polilla del tomate, esfíngide del algodón, esfíngide <i>Geometridae</i> y esfíngide de la baya silvestre; familia <i>Notodontidae</i> como la Puss Mot (<i>Cerura vinula</i>) y la polilla del gusano del manzano; familia de polillas tigre como la polilla tigre adornada, la Yellow Woolly Bear (<i>Spilosoma virginea</i>), polilla <i>Acraea</i> Mot, polilla tigre, polilla tigre manchada, Colón, polilla ligia y polilla <i>Rattleback</i> ; polillas de la familia <i>Ctenucha</i> como la polilla <i>Ctenucha</i> de Virginia; familia de los imantidos como la polilla gitana, polilla marcada blanca; familia de los noctuidos como la polilla de ocho manchas, la bruja negra, gusano falso medidor de la alfalfa, Sweetheart Underwing (<i>Catocala</i>), Locust Underwing (<i>Euparthenos</i> <i>noctua</i>) y polilla Hebrew.
Uvas	1 día	
Grupo de los pomos o melónidos incluyendo: Manzana, manzana silvestre, níspero, Mayhaw (<i>Crataegus</i>), pera, nashi y membrillo.	14 días	
Grupo de las drupas incluyendo: Albahorque, cereza (dulce y amarga), nectarina, melocotón, ciruela (incluyendo ciruela Chickasaw, ciruela silvestre y ciruela japonesa), plúot o ciruela albaricoque, ciruela pasa (fresca).	14 días	
Grupo de nueces de árboles incluyendo: Almendra, nuez de la haya, nuez amazónica, nuez blanca, anacardo, castaña, chinquapiñ (<i>Castanopsis</i>), avellana de Lambert, nuez del nogal americano, macadamia, pistacho, pistacho, nuez del nogal (negra e inglesa)	7 días	
Frutas tropicales incluyendo: Aguacate, zapote negro, canistel, zapote mamey, mango, papaya, chicozapote y camito	1 día	
ÁRBOLES ORNAMENTALES, ARBUSTOS, PLANTAS DE FLORES Y DE COBERTURA DE TERRENO Use el Sevin® Insect Killer Concentrate como tratamiento localizado para las infestaciones de plagas en árboles ornamentales, arbustos, plantas de flores y plantas de cobertura de terreno. Rocíelo hasta que el líquido corra procurando cubrir todos los lados de las hojas y el tronco. Para las plantas de cobertura de terreno, plantas de follaje, anuales y perennes, aplique un rocío uniforme. Repita el tratamiento si las plagas regresan, pero la reaplicación se debe limitar a no más de una vez cada siete días. Algunas plantas como las anuales pueden ser sensibles a la solución de rocío final. Se debe tratar un área limitada y se debe observar durante una semana antes de aplicar a toda la plantación.		
TRATAMIENTOS EN EL PERÍMETRO EXTERIOR Apliquelo hasta que el líquido corra como un rocío residual en la vegetación como los árboles y arbustos alrededor de las edificaciones y otras estructuras. Para las hormigas, aplíquelo en las trayectorias, alrededor de las puertas y ventanas y en otros lugares donde puedan encontrarse las hormigas. Para evitar que las plagas entren en las edificaciones, aplíquelo a una franja de suelo y vegetación de 5 a 10 pies (1.52 m a 3.04 m) de ancho alrededor y en áreas adyacentes a la edificación. Trate también los cimientos hasta una altura de 2 a 3 pies (0.60 m a 0.91 m) donde los insectos estén activos y puedan encontrar una forma de entrar. La superficie a tratar debe estar seca al momento de la aplicación.		
CÉSPEDES Aplique uniformemente el producto por cada área de 250 pies cuadrados (23.22 m ²). Para lograr mejores resultados, aplíquelo después de podar (el césped no debe ser mayor de 3 pulg. (7.62 cm) al momento de la aplicación). Repita la aplicación si es necesario solo si se observan señales de actividad renovada de los insectos.		
PLAGAS QUE CONTROLA		
Orugas, mariposas y polillas Orugas teje capullos tipo tiendas, orugas teje capullo y orugas que forman crisálidas. Orugas que comen hojas (incluyendo hojas de plantas florecientes, arbustos, otras plantas ornamentales, vegetales y árboles), minadores de la corteza, gusanos verdes, orugas que comen frutas, orugas que comen legumbres, orugas que comen vegetales, orugas barrenadoras de tallos y troncos, y orugas enrolladoras de hojas. Orugas de la familia de los prodoxidos como las polillas de la yuca; la familia de polillas de ropa, la familia del gusano de bolita, la familia de polillas Gelechiid como las polillas de la papa, la familia de polillas		
		Maquiledos comunes Familia de pectícelos de plata como el pectícelo de plata y el insecto de fuego Colémbolos Familia de los colémbolos como el colémbolo del bisalto y el colémbolo de la costa; colémbolos de la familia de los entomobríidos como los <i>Arsinoides</i> y globular Cucarachas Familia de los blátidos como la cucaracha oriental (cucaracha asiática o cucaracha Shad) y la cucaracha americana; familia de los blátidos como la cucaracha alemana (cucaracha rubia)

NOTE: The total number of pages must be divisible by four (4).

The maximum number of pages for this book is forty (40).

BOOKLET PAGES


BOOKLET TEXT (Pages 2-40)

.25"		4.25"		.25"																					
.25"				6.0"																					
5.5"																									
.25"		4.75"																							
<table><tr><th>PLAGAS QUE CONTROLA</th></tr><tr><td>Termitas</td></tr><tr><td>Familia de termitas que pudren la madera como la termita de la costa del Pacífico; termitas de la familia de termitas subterráneas. El producto no está concebido para reemplazar los tratamientos estructurales. Para las termitas —no se recomienda como protección exclusiva contra las termitas. Tales aplicaciones no sustituyen las alteraciones mecánicas ni el tratamiento del suelo o los cimientos sino son meramente un suplemento. Para las infestaciones activas, contrate una inspección profesional.</td></tr><tr><td>Tijeretas</td></tr><tr><td>Familia de tijeretas de antenas largas como el tijeretas de patas anilladas y el tijeretas ribereño; familia de tijeretas comunes como el tijeretas europeo</td></tr><tr><td>Saltamontes y grillos</td></tr><tr><td>Familia de grillos y saltamontes pigmeos como el saltamonte pigmeo azteca; familia de saltamontes de antenas cortas como el saltamontes del arbusto de la gobernadora, saltamontes Lubber, saltamontes pintado, saltamontes de Carolina, saltamontes dragón, Three-banded Grasshopper (Hadrotettix trifasciatus), saltamontes diferencial, langosta de patas rojas, saltamontes Melanoplus, saltamontes con manchas de pantera, saltamontes Lubber del sur, saltamontes del pájaro alutacea, saltamontes de pájaro americano, saltamontes del valle verde, Toothpick Grasshopper (Achurum carinatum), Horse Lubber Grasshopper (Taeniopoda eques), Pallid-winged Grasshopper (Trimerotropis pallidipennis) y Great Crested Grasshopper (Tropidolophus formosus); familia de los longicornios como el saltamontes longicornio, grillo de matorral, grillo mormón, Short-legged Shield back Katydid (Microcentrum californicum), grillo de California, Angularwinged Katydid (Microcentrum retinerve), Keeled Shield-back Katydid (Neduba carinata), grillo cabeza de cono, grillo ninfa de gladiador, grillo mescalero, grillo real, Fork-tailed Bush Katydid (Scudderella furcata); familia de grillo del camello como el grillo del camello manchado, grillo de la hueva, cara de niño; familia de los grillos como el grillo doméstico, grillo de campo, grillo del árbol de California, grillo del árbol nevado, grillo del árbol negro con cuernos; familia de los grillos topos como el grillo topo del norte y el grillo topo europeo.</td></tr><tr><td>Hemipteros</td></tr><tr><td>Familia de los miridos como la chinche Adelphocoris, chinche de planta escarlata, chinche de planta opaca, chinche de cuatro líneas; familia de los reduvidos o insectos asesinos como los asesinos de abejas, chinche Arilus cristatus, chinche Phymatidae, chinche Phymata; familia de los tingidos como la chinche del roble; familia de los pulgones como la ninfa pequeña del este, pulgón o áfido Myodocha y pulgón; familia de los berfidos como los pulgones berfidos; familia de las chinches con patas en forma de hoja como la chinche de Florida y la chinche hedionda; familia Niethrea louisianica como la chinche del este y la chinche del oeste; familia de las chinches hediondas como la chinche hedionda verde, chinche apesosa marrón mármol, chinche carnívora, chinche arlequín y chinche Podisus maculiventris</td></tr><tr><td>Cigarras y similares</td></tr><tr><td>Familia de las cigarras como las cigarras periódicas, la cigarra anual y la cigarra del oeste</td></tr><tr><td>Membranácidos</td></tr><tr><td>Familia de los membranácidos como el membranácido tipo espina, el membranácido del roble, el membranácido búfalo y el membranácido tipo langosta</td></tr><tr><td>Saltahojas</td></tr><tr><td>Familia de los saltahojas como el saltahoja de la vid, el saltahojas escaleta y verde y la chicharrita</td></tr><tr><td>Tisanópteros o trips</td></tr><tr><td>Familia de trips de franjas como el trip de alas de franjas; familia de los trips comunes</td></tr></table>				PLAGAS QUE CONTROLA	Termitas	Familia de termitas que pudren la madera como la termita de la costa del Pacífico; termitas de la familia de termitas subterráneas. El producto no está concebido para reemplazar los tratamientos estructurales. Para las termitas —no se recomienda como protección exclusiva contra las termitas. Tales aplicaciones no sustituyen las alteraciones mecánicas ni el tratamiento del suelo o los cimientos sino son meramente un suplemento. Para las infestaciones activas, contrate una inspección profesional.	Tijeretas	Familia de tijeretas de antenas largas como el tijeretas de patas anilladas y el tijeretas ribereño; familia de tijeretas comunes como el tijeretas europeo	Saltamontes y grillos	Familia de grillos y saltamontes pigmeos como el saltamonte pigmeo azteca; familia de saltamontes de antenas cortas como el saltamontes del arbusto de la gobernadora, saltamontes Lubber, saltamontes pintado, saltamontes de Carolina, saltamontes dragón, Three-banded Grasshopper (Hadrotettix trifasciatus), saltamontes diferencial, langosta de patas rojas, saltamontes Melanoplus, saltamontes con manchas de pantera, saltamontes Lubber del sur, saltamontes del pájaro alutacea, saltamontes de pájaro americano, saltamontes del valle verde, Toothpick Grasshopper (Achurum carinatum), Horse Lubber Grasshopper (Taeniopoda eques), Pallid-winged Grasshopper (Trimerotropis pallidipennis) y Great Crested Grasshopper (Tropidolophus formosus); familia de los longicornios como el saltamontes longicornio, grillo de matorral, grillo mormón, Short-legged Shield back Katydid (Microcentrum californicum), grillo de California, Angularwinged Katydid (Microcentrum retinerve), Keeled Shield-back Katydid (Neduba carinata), grillo cabeza de cono, grillo ninfa de gladiador, grillo mescalero, grillo real, Fork-tailed Bush Katydid (Scudderella furcata); familia de grillo del camello como el grillo del camello manchado, grillo de la hueva, cara de niño; familia de los grillos como el grillo doméstico, grillo de campo, grillo del árbol de California, grillo del árbol nevado, grillo del árbol negro con cuernos; familia de los grillos topos como el grillo topo del norte y el grillo topo europeo.	Hemipteros	Familia de los miridos como la chinche Adelphocoris, chinche de planta escarlata, chinche de planta opaca, chinche de cuatro líneas; familia de los reduvidos o insectos asesinos como los asesinos de abejas, chinche Arilus cristatus, chinche Phymatidae, chinche Phymata; familia de los tingidos como la chinche del roble; familia de los pulgones como la ninfa pequeña del este, pulgón o áfido Myodocha y pulgón; familia de los berfidos como los pulgones berfidos; familia de las chinches con patas en forma de hoja como la chinche de Florida y la chinche hedionda; familia Niethrea louisianica como la chinche del este y la chinche del oeste; familia de las chinches hediondas como la chinche hedionda verde, chinche apesosa marrón mármol, chinche carnívora, chinche arlequín y chinche Podisus maculiventris	Cigarras y similares	Familia de las cigarras como las cigarras periódicas, la cigarra anual y la cigarra del oeste	Membranácidos	Familia de los membranácidos como el membranácido tipo espina, el membranácido del roble, el membranácido búfalo y el membranácido tipo langosta	Saltahojas	Familia de los saltahojas como el saltahoja de la vid, el saltahojas escaleta y verde y la chicharrita	Tisanópteros o trips	Familia de trips de franjas como el trip de alas de franjas; familia de los trips comunes	<table><tr><th>PLAGAS QUE CONTROLA</th></tr><tr><td>Escarabajos</td></tr><tr><td>Familia de los escarabajos tigras como el escarabajo tigre hermoso, escarabajo tigre fino, escarabajo tigre verde, escarabajo tigre de noche; familia de los escarabajos de tierra como el escarabajo bombardero, escarabajo verde brillante, escarabajo de tierra europeo, escarabajo depredador verde, escarabajo de tierra común negro; familia de los escarabajos acuáticos como el escarabajo acuático plano, escarabajo buceador y escarabajo buceador marmoleado; familia de los grínidos como el escarabajo grínido grande y el escarabajo grínido pequeño; familia de los histéricos como los escarabajos depredadores de larvas; familia de los estafilínidos como el escarabajo marrón y dorado y el Thinopinus pictus; familia de los escarabajos enterradores o silfidos como el escarabajo enterrador, escarabajo enterrador de cuello dorado, escarabajo enterrador americano y escarabajo enterrador del norte; familia de lucánidos o ciervos volantes como el escarabajo ciervo de elefante, escarabajo de cuerno plano, escarabajo marrón rojizo y el escarabajo rugoso; familia de los pasálidos como el escarabajo de Jerusalén, familia de los escarabeidos como el escarabajo pelotero, escarabajo de orfebre, escarabajo de la vid, escarabajos Phyllophaga, escarabajo Polyphyla decemlineata y escarabajo japonés; familia de los buprestidos como el escarabajo buprestido dorado, barrenador de brote de pino del oeste, barrenador de cabeza plana del cerezo; familia de los eláteros, cascarudos o elatéridos como el escarabajo cascarudo de ojos grandes y escarabajos de fuego; familia de las luciérnagas como la luciérnaga de Pyralis y la luciérnaga de Pennsylvania; familia de los cantáridos o escarabajos soldado como el escarabajo soldado de Pennsylvania y el soldado escarabajo; familia de los escarabajos alas de red como el escarabajo alas en banda y el escarabajo de alas doradas; familia de los escarabajos arañas como el escarabajo araña y el escarabajo araña de Texas; familia de los escarabajos barrenadores de ramas como el barrenador de ramas del manzano; familia de los cléricos como el escarabajo clérico, el escarabajo clérico de California, escarabajo elegante y clérico rojo-azul; familia de los escarabajos planos como el escarabajo plano de la corteza; familia de escarabajos de los hongos o erotílidos como el escarabajo de hongo áspero; familia de los coccinélidos o mariquitas como la mariquita de dos manchas, mariquita de nueve manchas, mariquita Coccinella novemnotata, la mariquita Hippodamia convergens y la mariquita gris; familia de los tenebriónidos como el escarabajo de hongo con cuernos, tenebriónidos de cuello ancho, escarabajo plegado, escarabajo gusano de la harina y escarabajo Zopherinae; familia de los escarabajos de fuego o escarabajos color de fuego como el escarabajo de fuego; familia de los meloidos como el meloido rayado, el escarabajo meloido de Arizona y el escarabajo de alas cortas; familia de los cerambycids o escarabajos longicornios como el barrenador del pino o Callidium antennatum hesperum, barrenador del abeto Douglas, escarabajo longicornio, escarabajo Desmocerus palliatus, escarabajos podadores, chinche con cuello de espinas, langosta Megacyllene robiniae, chinche de manchas blancas, bórer Neocyttus acuminatus, barrenador de ramas de algodón, barrenador del algodón, barrenador de raíces gigante, barrenador del pino acanalado, barrenador del laurel de California, escarabajo eburia, escarabajo de las apocináceas rojo, escarabajo longicornio de flor y barrenador del sauce; familia de los barrenadores de semillas como los brúquidos o lánidos; familia de los escarabajos de hoja o crisomélidos como el escarabajo Calligrapha, escarabajo tortuga de algodoncillo, escarabajo de hoja de Dogbane, escarabajo de hoja de algodoncillo, escarabajo de la hoja del sauce, escarabajo manchado del espárrago, escarabajo tortuga claviforme, escarabajo manchado del espárrago del pepino, escarabajo de la hoja de lirio acuático, escarabajo de hoja de algodoncillo de pantano, escarabajo de la papa, escarabajo de la papa de Colorado o dorifora y escarabajo de hoja rodeado, familia de los gorgojos primitivos como el gorgojo primitivo del roble; familia de los picudos y familia de los gorgojos como el gorgojo del algodón, gorgojo de bellota de roble negro, gorgojo de hoja de fréjol menor, gorgojo de alfalfa, gorgojo del pino, gorgojo de la rosa, picudo del agave y gorgojos de los granos; familia de los Escolitinos o taladrillos como los escarabajos descortezadores del</td></tr></table>		PLAGAS QUE CONTROLA	Escarabajos	Familia de los escarabajos tigras como el escarabajo tigre hermoso, escarabajo tigre fino, escarabajo tigre verde, escarabajo tigre de noche; familia de los escarabajos de tierra como el escarabajo bombardero, escarabajo verde brillante, escarabajo de tierra europeo, escarabajo depredador verde, escarabajo de tierra común negro; familia de los escarabajos acuáticos como el escarabajo acuático plano, escarabajo buceador y escarabajo buceador marmoleado; familia de los grínidos como el escarabajo grínido grande y el escarabajo grínido pequeño; familia de los histéricos como los escarabajos depredadores de larvas; familia de los estafilínidos como el escarabajo marrón y dorado y el Thinopinus pictus; familia de los escarabajos enterradores o silfidos como el escarabajo enterrador, escarabajo enterrador de cuello dorado, escarabajo enterrador americano y escarabajo enterrador del norte; familia de lucánidos o ciervos volantes como el escarabajo ciervo de elefante, escarabajo de cuerno plano, escarabajo marrón rojizo y el escarabajo rugoso; familia de los pasálidos como el escarabajo de Jerusalén, familia de los escarabeidos como el escarabajo pelotero, escarabajo de orfebre, escarabajo de la vid, escarabajos Phyllophaga, escarabajo Polyphyla decemlineata y escarabajo japonés; familia de los buprestidos como el escarabajo buprestido dorado, barrenador de brote de pino del oeste, barrenador de cabeza plana del cerezo; familia de los eláteros, cascarudos o elatéridos como el escarabajo cascarudo de ojos grandes y escarabajos de fuego; familia de las luciérnagas como la luciérnaga de Pyralis y la luciérnaga de Pennsylvania; familia de los cantáridos o escarabajos soldado como el escarabajo soldado de Pennsylvania y el soldado escarabajo; familia de los escarabajos alas de red como el escarabajo alas en banda y el escarabajo de alas doradas; familia de los escarabajos arañas como el escarabajo araña y el escarabajo araña de Texas; familia de los escarabajos barrenadores de ramas como el barrenador de ramas del manzano; familia de los cléricos como el escarabajo clérico, el escarabajo clérico de California, escarabajo elegante y clérico rojo-azul; familia de los escarabajos planos como el escarabajo plano de la corteza; familia de escarabajos de los hongos o erotílidos como el escarabajo de hongo áspero; familia de los coccinélidos o mariquitas como la mariquita de dos manchas, mariquita de nueve manchas, mariquita Coccinella novemnotata, la mariquita Hippodamia convergens y la mariquita gris; familia de los tenebriónidos como el escarabajo de hongo con cuernos, tenebriónidos de cuello ancho, escarabajo plegado, escarabajo gusano de la harina y escarabajo Zopherinae; familia de los escarabajos de fuego o escarabajos color de fuego como el escarabajo de fuego; familia de los meloidos como el meloido rayado, el escarabajo meloido de Arizona y el escarabajo de alas cortas; familia de los cerambycids o escarabajos longicornios como el barrenador del pino o Callidium antennatum hesperum, barrenador del abeto Douglas, escarabajo longicornio, escarabajo Desmocerus palliatus, escarabajos podadores, chinche con cuello de espinas, langosta Megacyllene robiniae, chinche de manchas blancas, bórer Neocyttus acuminatus, barrenador de ramas de algodón, barrenador del algodón, barrenador de raíces gigante, barrenador del pino acanalado, barrenador del laurel de California, escarabajo eburia, escarabajo de las apocináceas rojo, escarabajo longicornio de flor y barrenador del sauce; familia de los barrenadores de semillas como los brúquidos o lánidos; familia de los escarabajos de hoja o crisomélidos como el escarabajo Calligrapha, escarabajo tortuga de algodoncillo, escarabajo de hoja de Dogbane, escarabajo de hoja de algodoncillo, escarabajo de la hoja del sauce, escarabajo manchado del espárrago, escarabajo tortuga claviforme, escarabajo manchado del espárrago del pepino, escarabajo de la hoja de lirio acuático, escarabajo de hoja de algodoncillo de pantano, escarabajo de la papa, escarabajo de la papa de Colorado o dorifora y escarabajo de hoja rodeado, familia de los gorgojos primitivos como el gorgojo primitivo del roble; familia de los picudos y familia de los gorgojos como el gorgojo del algodón, gorgojo de bellota de roble negro, gorgojo de hoja de fréjol menor, gorgojo de alfalfa, gorgojo del pino, gorgojo de la rosa, picudo del agave y gorgojos de los granos; familia de los Escolitinos o taladrillos como los escarabajos descortezadores del
PLAGAS QUE CONTROLA																									
Termitas																									
Familia de termitas que pudren la madera como la termita de la costa del Pacífico; termitas de la familia de termitas subterráneas. El producto no está concebido para reemplazar los tratamientos estructurales. Para las termitas —no se recomienda como protección exclusiva contra las termitas. Tales aplicaciones no sustituyen las alteraciones mecánicas ni el tratamiento del suelo o los cimientos sino son meramente un suplemento. Para las infestaciones activas, contrate una inspección profesional.																									
Tijeretas																									
Familia de tijeretas de antenas largas como el tijeretas de patas anilladas y el tijeretas ribereño; familia de tijeretas comunes como el tijeretas europeo																									
Saltamontes y grillos																									
Familia de grillos y saltamontes pigmeos como el saltamonte pigmeo azteca; familia de saltamontes de antenas cortas como el saltamontes del arbusto de la gobernadora, saltamontes Lubber, saltamontes pintado, saltamontes de Carolina, saltamontes dragón, Three-banded Grasshopper (Hadrotettix trifasciatus), saltamontes diferencial, langosta de patas rojas, saltamontes Melanoplus, saltamontes con manchas de pantera, saltamontes Lubber del sur, saltamontes del pájaro alutacea, saltamontes de pájaro americano, saltamontes del valle verde, Toothpick Grasshopper (Achurum carinatum), Horse Lubber Grasshopper (Taeniopoda eques), Pallid-winged Grasshopper (Trimerotropis pallidipennis) y Great Crested Grasshopper (Tropidolophus formosus); familia de los longicornios como el saltamontes longicornio, grillo de matorral, grillo mormón, Short-legged Shield back Katydid (Microcentrum californicum), grillo de California, Angularwinged Katydid (Microcentrum retinerve), Keeled Shield-back Katydid (Neduba carinata), grillo cabeza de cono, grillo ninfa de gladiador, grillo mescalero, grillo real, Fork-tailed Bush Katydid (Scudderella furcata); familia de grillo del camello como el grillo del camello manchado, grillo de la hueva, cara de niño; familia de los grillos como el grillo doméstico, grillo de campo, grillo del árbol de California, grillo del árbol nevado, grillo del árbol negro con cuernos; familia de los grillos topos como el grillo topo del norte y el grillo topo europeo.																									
Hemipteros																									
Familia de los miridos como la chinche Adelphocoris, chinche de planta escarlata, chinche de planta opaca, chinche de cuatro líneas; familia de los reduvidos o insectos asesinos como los asesinos de abejas, chinche Arilus cristatus, chinche Phymatidae, chinche Phymata; familia de los tingidos como la chinche del roble; familia de los pulgones como la ninfa pequeña del este, pulgón o áfido Myodocha y pulgón; familia de los berfidos como los pulgones berfidos; familia de las chinches con patas en forma de hoja como la chinche de Florida y la chinche hedionda; familia Niethrea louisianica como la chinche del este y la chinche del oeste; familia de las chinches hediondas como la chinche hedionda verde, chinche apesosa marrón mármol, chinche carnívora, chinche arlequín y chinche Podisus maculiventris																									
Cigarras y similares																									
Familia de las cigarras como las cigarras periódicas, la cigarra anual y la cigarra del oeste																									
Membranácidos																									
Familia de los membranácidos como el membranácido tipo espina, el membranácido del roble, el membranácido búfalo y el membranácido tipo langosta																									
Saltahojas																									
Familia de los saltahojas como el saltahoja de la vid, el saltahojas escaleta y verde y la chicharrita																									
Tisanópteros o trips																									
Familia de trips de franjas como el trip de alas de franjas; familia de los trips comunes																									
PLAGAS QUE CONTROLA																									
Escarabajos																									
Familia de los escarabajos tigras como el escarabajo tigre hermoso, escarabajo tigre fino, escarabajo tigre verde, escarabajo tigre de noche; familia de los escarabajos de tierra como el escarabajo bombardero, escarabajo verde brillante, escarabajo de tierra europeo, escarabajo depredador verde, escarabajo de tierra común negro; familia de los escarabajos acuáticos como el escarabajo acuático plano, escarabajo buceador y escarabajo buceador marmoleado; familia de los grínidos como el escarabajo grínido grande y el escarabajo grínido pequeño; familia de los histéricos como los escarabajos depredadores de larvas; familia de los estafilínidos como el escarabajo marrón y dorado y el Thinopinus pictus; familia de los escarabajos enterradores o silfidos como el escarabajo enterrador, escarabajo enterrador de cuello dorado, escarabajo enterrador americano y escarabajo enterrador del norte; familia de lucánidos o ciervos volantes como el escarabajo ciervo de elefante, escarabajo de cuerno plano, escarabajo marrón rojizo y el escarabajo rugoso; familia de los pasálidos como el escarabajo de Jerusalén, familia de los escarabeidos como el escarabajo pelotero, escarabajo de orfebre, escarabajo de la vid, escarabajos Phyllophaga, escarabajo Polyphyla decemlineata y escarabajo japonés; familia de los buprestidos como el escarabajo buprestido dorado, barrenador de brote de pino del oeste, barrenador de cabeza plana del cerezo; familia de los eláteros, cascarudos o elatéridos como el escarabajo cascarudo de ojos grandes y escarabajos de fuego; familia de las luciérnagas como la luciérnaga de Pyralis y la luciérnaga de Pennsylvania; familia de los cantáridos o escarabajos soldado como el escarabajo soldado de Pennsylvania y el soldado escarabajo; familia de los escarabajos alas de red como el escarabajo alas en banda y el escarabajo de alas doradas; familia de los escarabajos arañas como el escarabajo araña y el escarabajo araña de Texas; familia de los escarabajos barrenadores de ramas como el barrenador de ramas del manzano; familia de los cléricos como el escarabajo clérico, el escarabajo clérico de California, escarabajo elegante y clérico rojo-azul; familia de los escarabajos planos como el escarabajo plano de la corteza; familia de escarabajos de los hongos o erotílidos como el escarabajo de hongo áspero; familia de los coccinélidos o mariquitas como la mariquita de dos manchas, mariquita de nueve manchas, mariquita Coccinella novemnotata, la mariquita Hippodamia convergens y la mariquita gris; familia de los tenebriónidos como el escarabajo de hongo con cuernos, tenebriónidos de cuello ancho, escarabajo plegado, escarabajo gusano de la harina y escarabajo Zopherinae; familia de los escarabajos de fuego o escarabajos color de fuego como el escarabajo de fuego; familia de los meloidos como el meloido rayado, el escarabajo meloido de Arizona y el escarabajo de alas cortas; familia de los cerambycids o escarabajos longicornios como el barrenador del pino o Callidium antennatum hesperum, barrenador del abeto Douglas, escarabajo longicornio, escarabajo Desmocerus palliatus, escarabajos podadores, chinche con cuello de espinas, langosta Megacyllene robiniae, chinche de manchas blancas, bórer Neocyttus acuminatus, barrenador de ramas de algodón, barrenador del algodón, barrenador de raíces gigante, barrenador del pino acanalado, barrenador del laurel de California, escarabajo eburia, escarabajo de las apocináceas rojo, escarabajo longicornio de flor y barrenador del sauce; familia de los barrenadores de semillas como los brúquidos o lánidos; familia de los escarabajos de hoja o crisomélidos como el escarabajo Calligrapha, escarabajo tortuga de algodoncillo, escarabajo de hoja de Dogbane, escarabajo de hoja de algodoncillo, escarabajo de la hoja del sauce, escarabajo manchado del espárrago, escarabajo tortuga claviforme, escarabajo manchado del espárrago del pepino, escarabajo de la hoja de lirio acuático, escarabajo de hoja de algodoncillo de pantano, escarabajo de la papa, escarabajo de la papa de Colorado o dorifora y escarabajo de hoja rodeado, familia de los gorgojos primitivos como el gorgojo primitivo del roble; familia de los picudos y familia de los gorgojos como el gorgojo del algodón, gorgojo de bellota de roble negro, gorgojo de hoja de fréjol menor, gorgojo de alfalfa, gorgojo del pino, gorgojo de la rosa, picudo del agave y gorgojos de los granos; familia de los Escolitinos o taladrillos como los escarabajos descortezadores del																									

NOTE: The total number of pages must be divisible by four (4).
The maximum number of pages for this book is forty (40).

BOOKLET PAGES

BOOKLET TEXT (Pages 2-40)



NOTE: The total number of pages must be divisible by four (4).

The maximum number of pages for this book is forty (40).

BOOKLET PAGES

BOOKLET TEXT (Pages 2-40)

.25"		4.25"		.25"	
.25"		5.5"		6.0"	
.25"		4.75"		14	
Plantas	Plagas	Uso antes de la cosecha	Plantas	Plagas	Uso antes de la cosecha
VEGETALES			VEGETALES		
• Cilantro • Canónigo • Berro o mastuerzo • Berro de montaña • Diente de león • Romaza o vinagrera • Endivia • Hinojo de Florencia • Lechuga de cogollo (beluga) • Lechuga • Abanico • Perejil • Verdolaga (de jardín) • Verdolaga (de invierno) • Radichio (achicoria roja) • Ruibarbo • Espinaca • Acelga	de la cebolla, chinche hedionda y cascarudo o saltaperico		• Repollo chino (napa) • Hortalizas Brassica de hojas comestibles incluyendo: grelo (brócoli rabe o raab); • Repollo chino (bok choy) • Mostaza china (gai choy) • Berza • Col rizada • Nipposinica o mostaza japonesa • Mostaza de hoja • Espinaca japonesa • Raps o colza	saltaperico (adultos), áfido y mosca blanca	
Vegetales que dan frutos incluyendo: • Berenjena • Uchuva (<i>Physalis</i>) • Pepino melón (pepino dulce o melón de árbol) • Pimiento (incluyendo el pimiento o ají, pimiento chileno, pimiento de cocinar, pimiento y pimiento dulce) • Tomatillo • Tomate	Gusano cogollero, gusano cogollero del sur, gusano cogollero real, Yellowstriped Celery Leaf Tier (<i>Udea rubigalis</i>), escarabajo de la papa de Colorado, bórer del maíz europeo, bórer del maíz sureño, escarabajo del pepino, gusano cortador, alticino, gusano tejedor de jardín, chinche hedionda verde, gusano, minador de hojas (adultos), saltahoja, cercopídeo de campo, gusano del pimiento (adultos), gorgojo del pimiento, polilla de las plantas, gusano de la yema del tabaco, gusano de la fruta del tomate, gusano del tomate (<i>Keiferia lycopersicella</i>), áfidos, gusano cogollero, gusano cogollero de la remolacha, gusano falso medidor de la col de otoño, saltamonte, chinche opaca, chinche hedionda marrón, psílido o piojo saltarín del tomate, trip y mosca blanca.	Rocielo hasta 1 día antes de la cosecha.	Vegetales de raíces y tubérculos • Arracacha • Maranta • Alcachofa • Remolacha de huerto • Lampazo • Platanillo comestible • Zanahoria • Vegetal de la raíz del perifollo • Achicoria • Chufa • Malanga (Taro) • Jengibre • Ginseng • Rábano picante • Lerén • Vegetal de la raíz del perejil • Chirivía • Papa • Rábano blanco o rábano japonés (daikon) • Colinabo • Salsifi (barba cabruna) • Escorzonera negra • Escorzonera • Escaravia • Camote • Yautía • Cúrcuma • Nabo • Jicama • Ñame	Gusano cortador, gusano falso medidor de la col, escarabajo del pepino, bórer del maíz europeo, alticino, gusano de la raíz del maíz sureño (adulto), gorgojo o picudo de los vegetales, escarabajo Naupactus (adulto), gusano cogollero, gusano cogollero de la remolacha, gusano de franjas amarillas de la col, escarabajo de la papa de Colorado, saltamonte, gusano de la col importado, saltahoja de la papa, psílido de la papa, gusano de la papa y chinche de las plantas opaca	Rocielo hasta 1 día antes de la cosecha.
Vegetales del género Brassica incluyendo: • Brócoli • Brócoli chino (gai lan, brócoli blanco floreciente) • Col de Bruselas • Coliflor • Romanesco • Colirrábano • Repollo	Gusano cogollero del maíz, escarabajo del pepino, gusano cortador, palomilla dorso de diamante, alticino, gusano de la col importado, saltahoja, gusano peludo, gusano de la col sureño, gusano de la yema del tabaco, gusano falso medidor de la alfalfa, gusano cogollero, gusano falso medidor de la col, gusano tejedor de la col, escarabajo de tierra, minador de hojas (adultos), chinche opaca, trip de la cebolla, chinche hedionda, cascarudo o	Rocielo hasta 1 día antes de la cosecha.			

**NOTE: The total number of pages must be divisible by four (4).
The maximum number of pages for this book is forty (40).**

BOOKLET PAGES

BOOKLET TEXT (Pages 2-40)

.25"			4.25"			.25"		
.25"			5.5"			6.0"		
.25"								
Plantas	Plagas	Uso antes de la cosecha	Plantas	Plagas	Uso antes de la cosecha			
VEGETALES			VEGETALES					
Vegetales cucurbitáceos incluyendo, pero sin limitarse a:	Gusano cortador, gusano falso medidor de la col, escarabajo del pepino (adulto), saltahoja, gusano del melón, gusano del pepino, gusano del melón, gusano de la calabaza, bórer de la enredadera de calabaza, gusano cogollero del maíz, polilla de las plantas y chinche hedionda	Rocielo hasta 1 día antes de la cosecha.	• Sandía (incluyendo los híbridos y variedades).					
• Chayote (fruta)			Vegetales de bulbos (especies de Allium) incluyendo:	Gusano cogollero, gusano cortador, minador de hojas (adultos), gusano de la cebolla (adultos), chinche hedionda, áfido y trip de la cebolla	Rocielo hasta 7 días antes de la cosecha.			
• Calabaza china			• Ajo; ajo elefante					
• Melón Kalahari			• Chalota verde o echalote verde					
• Pepino			• Cebolleta o cebolleta					
• Pepinillo			• Puerro o ajo porro					
• Calabaza (comestible, incluyendo jicaro o acocote, cucuzza, luffa o estropajo y ocrá china)			• Cebolla verde o cebollino, cebolla de bulbo seco					
• Variedades de Momordica (incluyendo balsam apple (Momordica balsamina), melón amargo o cundeamor chino, melón amargo y pepino chino)			• Cebolleta o cebolla larga					
• Melón (híbridos y/o variedades de Cucumis melo) (incluye el cantalupo genuino, cantalupo, melón verde o casaba, melón Crenshaw, melón Golden Pershaw, melón dulce, honey balls, mango melon (Cucumis melo var. chito), melón persa, melón Kalahari, melón piel de sapo y melón serpiente o pepino serpiente)			• Chalotas de bulbo seco y verde					
• Calabaza			• Cebolla de verdeo, cebollino o cebolleta					
• Calabaza de verano (incluyendo crookneck squash (Cucurbita pepo), calabacín, calabaza de verano amarilla, calabacita y calabacín)			Girasol	Gusano cortador, gusano blanco y cascarudo o saltaperico	Rocielo hasta 30 días antes de la cosecha.			
• Calabaza de invierno (incluyendo zapallo, calabaza, calabaza gigante, calabaza de Des Moines y calabaza común)			Maíz dulce	Chinche, gusano de la raíz del maíz (adulto), mosca del maíz (Euxesta stigmatias), gusano cortador, alticino, saltahoja, escarabajo japonés (adulto), escarabajo nitidulido (adultos), chinche de las plantas opaca, gusano cogollero, bórer del maíz, gusano cogollero del maíz, saltamonte y áfido	Rocielo hasta 3 días antes de la cosecha.			
			Legumbres — jugosas y secas	Gusano cortador, oruga de la dama pintada, gusano peludo, Silverspotted Skipp (Epargyreus clarus), oruga de la alfalfa, gusano cogollero, gusano cogollero del sur, escarabajo de la hoja del frijol de franjas amarillas, escarabajo de la papa de Colorado, bórer del maíz europeo, bórer del maíz, gusano cogollero del maíz sureño, escarabajo de la raíz del maíz (adulto), Curculio del frijol, escarabajo del pepino, alticino, gusano verde, escarabajo de tierra, gusano de la col importado, escarabajo japonés, devorador de hojas, saltahoja, minador de las hojas (adulto), escarabajo del frijol mexicano, gorgojo del guisante, gorgojo de las hojas del guisante, polilla de las plantas, saltahoja de la papa, escarabajo del grano de maíz, gusano del grano del maíz (adulto),	Rocielo hasta 1 día antes de la cosecha para los guisantes o frijoles jugosos sin cáscara o en vainas y agregue 21 días antes de la cosecha para los guisantes o frijoles sin cáscara secos			
			Guisantes de vainas comestibles					
			jugosos, guisantes sin cáscara					
			jugosos y guisantes sin cáscara secos (variedades de Pisum) incluyendo:					
			• Guisante enano					
			• Guisante de vaina comestible					
			• Guisante mollar					
			• Tirabeque dulce					
			• Gandul					
			• Petipuas					
			• Arveja					
			• Chicharo verde					
			• Lenteja					
			Frijoles de vainas comestibles					
			jugosos, frijoles sin					
					15			
.25"			4.75"			.25"		

NOTE: The total number of pages must be divisible by four (4).

The maximum number of pages for this book is forty (40).

BOOKLET TEXT
(Pages 2-40)

NOTE: The total number of pages must be divisible by four (4).
The maximum number of pages for this book is forty (40).

BOOKLET TEXT

(Pages 2-40)

NOTE: The total number of pages must be divisible by four (4).
The maximum number of pages for this book is forty (40).